

FCS
Math: Functions
Exercises

Massimo Caboara

April 29th, 2021

Solved exercise


Example 1. *We have the function*

$$F: \mathbb{R} \longrightarrow \mathbb{R}$$
$$x \longmapsto \begin{cases} \arctan(x) & x < 0 \\ x^2 + 1 & x \geq 0 \end{cases}$$

1. *Draw the graph of F . Mark the intersections with the axis and the interesting points.*
2. *Is F injective, surjective, invertible? What is the image of F ?*
3. *Determine $F(0)$, $F(\pi)$, $F([-\pi/2, \pi/4])$.*
4. *Determine $F^{-1}(0)$, $F^{-1}(1)$, $F^{-1}((-\sqrt{2}/2, \sqrt{2}/2])$, $F^{-1}((-\infty, 0])$.*
5. *Build an invertible function from F by restricting its domain and/or codomain.*
6. *Determine the formula for this inverse.*

1 *The graph is*

GNU26: Graph of $F(x)$



2 It is easy to see that every horizontal line $y = a$, $a \in \mathbb{R}$ intersects the graph of F once or not at all, so F is injective.

The horizontal line $y = 1/2$ does not intersect the graph of F , so F is not surjective. We could have chosen another horizontal line with no intersection, for example $y = -2$. The graphical rule says that a function is surjective if EVERY horizontal line $y = a$ with a in the domain of F intersects the graph of the function at least once.

Being injective but not surjective, F is not invertible.

The image of F is $(-\pi/2, 0] \cup [1, +\infty)$

3 $F(0) = 1$, $F([1, 2]) = [F(1), F(2)] = [2, 5]$ because F is increasing. To determine the image of $(-\pi/2, \pi/4]$ it is useful to split this interval

$$(-\pi/2, \pi/4] = (-\pi/2, 0) \cup [0, \pi/4]$$

and we have

$$\begin{aligned} F((-\pi/2, \pi/4]) &= F((-\pi/2, 0) \cup [0, \pi/4]) \\ &= F((-\pi/2, 0)) \cup F([0, \pi/4]) \\ &= (F(-\pi/2), F(0)) \cup [F(0), F(\pi/4)] \\ &= (-1, 0) \cup [1, \pi^2/16 + 1] \end{aligned}$$

4 $F^{-1}(1/2) = \emptyset$ because the graph of F and the line $y = 1/2$ have no intersection.

$F^{-1}(-2) = \emptyset$ because the graph of F and the line $y = -2$ have no intersection.

For the counterimage of the interval $[-1, 2]$ it is useful to split this interval

$$[-1, 2] = [-1, 0) \cup [0, 1) \cup [1, 2]$$

and we have

$$\begin{aligned}
 F^{-1}([-1, 2]) &= F^{-1}([-1, 0] \cup [0, 1] \cup [1, 2]) \\
 &= F^{-1}([-1, 0]) \cup F^{-1}([0, 1]) \cup F^{-1}([1, 2]) \\
 &= [F^{-1}(-1), F^{-1}(0)] \cup \emptyset \cup [F^{-1}(1), F^{-1}(2)] \\
 &= [-\pi/4, 0] \cup \emptyset \cup [0, 1] \\
 &= [-\pi/4, 0] \cup [0, 1] \\
 &= [-\pi/4, 1]
 \end{aligned}$$

5 The function is injective, so we keep its domain. To make the function surjective, we restrict the codomain to the image. The new, invertible, function is

$$\begin{aligned}
 F' : \mathbb{R} &\longrightarrow (-\pi/2, 0] \cup [1, +\infty) \\
 x &\mapsto \begin{cases} \arctan(x) & x < 0 \\ x^2 + 1 & x \geq 0 \end{cases}
 \end{aligned}$$

We compute its inverse piece by piece

First piece

$$\arctan(x) = y \iff \tan(\arctan(x)) = \tan(y) \iff x = \tan(y)$$

and we can apply the tangent to the equation because $\arctan(x) \in (-\pi/2, \pi/2)$ for every x and $y \in (-\pi/2, 0]$ for this piece. In both cases the values are in the domain of THIS tangent (restriction of the tangent to $(-\pi/2, \pi/2)$), that for the sake of simplicity we still call tangent

$$\begin{aligned}
 \tan : (-\pi/2, \pi/2) &\longrightarrow \mathbb{R} \\
 x &\mapsto \tan(x)
 \end{aligned}$$

and so we can apply it to the equation. Since THIS tangent is invertible, the equation does not change and we have $\tan(\arctan(x)) = x$

Second piece

$$x^2 + 1 = y \iff x^2 = y - 1 \iff \sqrt{x^2} = \sqrt{y - 1} \iff x = \sqrt{y - 1}$$

and we can apply the square root to the equation because $x^2 \geq 0$ and $y - 1 \geq 0$ in this piece. In both cases the values are in the domain of the square root, and so we can apply it to the equation. Since the square root

$$\begin{aligned}
 \sqrt{\cdot} : \mathbb{R}_0^+ &\longrightarrow \mathbb{R}_0^+ \\
 x &\mapsto \sqrt{x}
 \end{aligned}$$

is invertible with inverse

$$\begin{aligned}
 (\cdot)^2 : \mathbb{R}_0^+ &\longrightarrow \mathbb{R}_0^+ \\
 x &\mapsto x^2
 \end{aligned}$$

the equation does not change and we have $\sqrt{x^2} = x$ for the x 's in \mathbb{R}_0^+ .

The inverse of F' is then

$$F'^{-1} : (-\pi/2, 0] \cup [1, +\infty) \longrightarrow \mathbb{R}$$

$$x \longmapsto \begin{cases} \tan(x) & x \in (-\pi/2, 0] \\ x^2 + 1 & x \in [1, +\infty) \end{cases}$$

Proposed exercises

Exercise 1. We have the function

$$F : \mathbb{R} \longrightarrow \mathbb{R}$$

$$x \longmapsto \arctan x$$

1. Draw the graph of F . Mark the intersections with the axis and the interesting points.
2. Is F injective, surjective, invertible?
3. Determine $F(0)$, $F(\pi)$, $F([-\pi/2, \pi/4])$.
4. Determine $F^{-1}(0)$, $F^{-1}(1)$, $F^{-1}((-\sqrt{2}/2, \sqrt{2}/2])$, $F^{-1}((-\infty, 0])$.
5. Build an invertible function from F by restricting its domain and/or codomain.
6. Determine the formula for this inverse and draw its graph.

Exercise 2. We have the function

$$F : \mathbb{R} \longrightarrow \mathbb{R}$$

$$x \longmapsto \arctan 2^x$$

1. Draw the graph of F . Mark the intersections with the axis and the interesting points.
2. Is F injective, surjective, invertible?
3. Determine $F(0)$, $F(-1)$, $F([-\pi/2, \pi/4])$.
4. Determine $F^{-1}(0)$, $F^{-1}(1)$, $F^{-1}((0, 1])$, $F^{-1}((-\infty, 0])$.
5. Build an invertible function from F by restricting its domain and/or codomain.
6. Determine the formula for this inverse and draw its graph.

Exercise 3. We have the function

$$F : \mathbb{R} \longrightarrow \mathbb{R}$$

$$x \longmapsto 2^{\arctan x}$$

1. Draw the graph of F . Mark the intersections with the axis and the interesting points.
2. Is F injective, surjective, invertible?
3. Determine $F(0)$, $F(\pi)$, $F([-\pi/2, \pi/4])$.
4. Determine $F^{-1}(0)$, $F^{-1}(1)$, $F^{-1}((-\sqrt{2}/2, \sqrt{2}/2])$, $F^{-1}((-\infty, 0])$.
5. Build an invertible function from F by restricting its domain and/or codomain.
6. Determine the formula for this inverse and draw its graph.

Exercise 4. We have the function

$$F : \mathbb{R} \longrightarrow \mathbb{R}$$

$$x \longmapsto \begin{cases} 3^x & x < 0 \\ -x^2 & x \geq 0 \end{cases}$$

1. Draw the graph of F . Mark the intersections with the axis and the interesting points.
2. Is F injective, surjective, invertible?
3. Determine $F(0)$, $F(-1)$, $F([-1, 1])$.
4. Determine $F^{-1}(0)$, $F^{-1}(-1)$, $F^{-1}([-2, 2])$
5. Build an invertible function from F by restricting its domain and/or codomain.
6. Determine the formula for this inverse and draw its graph.

Exercise 5. We have the function

$$F : \mathbb{R} \longrightarrow \mathbb{R}$$

$$x \longmapsto \begin{cases} \sin(x) - 1 & x < 0 \\ \sqrt{x} & x \geq 0 \end{cases}$$

1. Draw the graph of F . Mark the intersections with the axis and the interesting points.
2. Is F injective, surjective, invertible?
3. Determine $F(0)$, $F(-\pi)$, $F([-\pi/2, 2])$.
4. Determine $F^{-1}(0)$, $F^{-1}(1)$, $F^{-1}([-1, 1])$.
5. Build an invertible function from F by restricting its domain and/or codomain.
6. Determine the formula for this inverse and draw its graph.