

# An ultrafilter approach to Jin's Theorem

Mathias Beiglböck\*

This paper is dedicated to Dona Strauss on the occasion of her 75th birthday.

## Abstract

It is well known and not difficult to prove that if  $C \subseteq \mathbb{Z}$  has positive upper Banach density, the set of differences  $C - C$  is syndetic, i.e. the length of gaps is uniformly bounded. More surprisingly, Renling Jin showed that whenever  $A$  and  $B$  have positive upper Banach density, then  $A - B$  is *piecewise syndetic*.

Jin's result follows trivially from the first statement provided that  $B$  has large intersection with a shifted copy  $A - n$  of  $A$ . Of course this will not happen in general if we consider shifts by integers, but the idea can be put to work if we allow "shifts by ultrafilters". As a consequence we obtain Jin's Theorem.

The *upper Banach density* of  $C \subseteq \mathbb{Z}$  is given by

$$d^*(C) := \overline{\lim}_{m \rightarrow \infty} \frac{|C \cap \{n, \dots, m\}|}{m - n + 1}.$$

A set  $S \subseteq \mathbb{Z}$  is syndetic if the gaps of  $S$  are of uniformly bounded length, i.e. if there exists some  $k > 0$  such that  $S - \{-k, \dots, k\} = \mathbb{Z}$ . A set  $P \subseteq \mathbb{Z}$  is piecewise syndetic if it is syndetic on large pieces, i.e. if there exists some  $k \geq 0$  such that  $P - \{-k, \dots, k\}$  contains arbitrarily long intervals of integers.

It was first noted by Følner ([Føl54a, Føl54b]) that  $C - C = \{c_1 - c_2 : c_1, c_2 \in C\}$  is syndetic provided that  $d^*(C) > 0$ . (To see this, pick a subset  $\{i_1, \dots, i_m\} \subseteq \mathbb{Z}$  which is *maximal* subject to the condition that  $C - i_1, \dots, C - i_m$  are mutually disjoint. This is possible since disjointness implies  $d^*(C - i_1 \cup \dots \cup C - i_m) = m \cdot d^*(C)$ , therefore  $m$  is at most  $1/d^*(C)$ . But then maximality implies that for each  $n \in \mathbb{Z}$  there is some  $i_k, k \in \{1, \dots, m\}$  such that  $(C - n) \cap (C - i_k) \neq \emptyset$ , resp.  $n \in (C - C) + i_k$ . Thus  $\bigcup_{k=1}^m (C - C) + i_k = \mathbb{Z}$ .)

Simple counterexamples yield that the analogous statement fails when two different sets are considered, but Renling Jin discovered the following interesting result.

**Theorem 1** ([Jin02]). *Let  $A, B \subseteq \mathbb{Z}, d^*(A), d^*(B) > 0$ . Then  $A - B$  is piecewise syndetic.*

In Section 1 we reprove Jin's Theorem by reducing it to the  $C - C$  case. Subsequently we discuss some modifications of our argument which allow to recover the refinements resp. generalizations of Jin's result found in [JK03, BFW06, Jin08, BBF].

---

\*2000 Mathematics Subject Classification. 11B05, 05D10

keywords and phrases. density, piecewise syndetic, piecewise Bohr, sumsets, ultrafilters

The author acknowledges financial support from the Austrian Science Fund (FWF) under grant P21209.

## 1 Ultrafilter proof of Jin's Theorem

As indicated in the abstract, we aim to show that one can shift a given large set  $A \subseteq \mathbb{Z}$  by an ultrafilter so that it will have large intersection with another, previously specified set. We motivate the definition of this ultrafilter-shift by means of analogy: For  $n \in \mathbb{Z}$  denote by  $e(n)$  the principle ultrafilter on  $\mathbb{Z}$  which corresponds to  $n$  and notice that

$$A - n = \{k \in \mathbb{Z} : n \in A - k\} = \{k \in \mathbb{Z} : A - k \in e(n)\}.$$

Given an ultrafilter  $p$  on  $\mathbb{Z}$ , we thus define  $\{k \in \mathbb{Z} : A - k \in p\}$  as the official meaning of “ $A - p$ ”.

**Lemma 2.** *Let  $A, B \subseteq \mathbb{Z}$ . Then there exists an ultrafilter  $p$  on  $\mathbb{Z}$  such that*

$$d^*((A - p) \cap B) = d^*(\{k : A - k \in p\} \cap B) \geq d^*(A) \cdot d^*(B).$$

The proof of Lemma 2 requires some preliminaries.

For a fixed set  $A \subseteq \mathbb{Z}$  we can choose an invariant mean, i.e. a shift invariant finitely additive probability measure  $\mu$  on  $(\mathbb{Z}, \mathcal{P}(\mathbb{Z}))$  such that  $d^*(A) = \mu(A)$ . This is well known among aficionados (see for instance [Ber06, Theorem 5.8]) and not difficult to prove. First pick a sequence of finite intervals  $I_n \subseteq \mathbb{Z}$  such that  $|I_n| \rightarrow \infty$  and  $d^*(A) = \lim_{n \rightarrow \infty} \mu_n(A)$  where  $\mu_n(B) := \frac{1}{|I_n|} |B \cap I_n|$  for  $B \in \mathcal{P}(B)$ . Then let  $\mu$  be a cluster point of the set  $\{\mu_n : n \in \mathbb{N}\}$  in the (compact) product topology of  $[0, 1]^{\mathcal{P}(\mathbb{Z})}$ .

We consider the Stone-Ćech compactification  $\beta\mathbb{Z}$  of the discrete space  $\mathbb{Z}$ . For our purpose it is convenient to view  $\beta\mathbb{Z}$  as the set of all ultrafilters on  $\mathbb{Z}$ . By identifying integers with principal ultrafilters,  $\mathbb{Z}$  is naturally embedded in  $\beta\mathbb{Z}$ . A clopen basis for the topology is given by the sets  $\overline{C} := \{p \in \beta\mathbb{Z} : C \in p\}$ , where  $C \subseteq \mathbb{Z}$ . A mean  $\mu$  on  $\mathbb{Z}$  gives rise to a positive linear functional  $\Lambda$  on the space  $B(\mathbb{Z})$  of bounded functions on  $\mathbb{Z}$ . Making the identification  $B(\mathbb{Z}) \cong C(\beta\mathbb{Z})$  we find that, by the Riesz representation Theorem, there exists a regular Borel probability measure  $\tilde{\mu}$  on  $\beta\mathbb{Z}$  which corresponds to the mean  $\mu$  in the sense that  $\mu(A) = \tilde{\mu}(\overline{A})$  for all  $A \subseteq \mathbb{Z}$ . (This procedure is carried out in detail for instance in [Pat88, p 11].)

*Proof of Lemma 2.* Pick a sequence of intervals  $I_n \subseteq \mathbb{Z}, |I_n| \uparrow \infty$  such that  $d^*(B) = \lim_{n \rightarrow \infty} \frac{|I_n \cap B|}{|I_n|}$ . Pick an invariant mean  $\mu$  such that  $\mu(A) = d^*(A)$ . Define  $f_n : \beta\mathbb{Z} \rightarrow [0, 1]$  by

$$f_n(p) := \frac{1}{|I_n|} \sum_{k \in I_n \cap B} \mathbb{1}_{\overline{A-k}}(p) = \frac{|I_n \cap B \cap \{k : A - k \in p\}|}{|I_n|}$$

and set  $f(p) := \overline{\lim}_{n \rightarrow \infty} f_n(p) \leq d^*(B \cap \{k : A - k \in p\})$ . By Fatou's Lemma

$$\int f d\tilde{\mu} \geq \overline{\lim}_{n \rightarrow \infty} \int \frac{1}{|I_n|} \sum_{k \in I_n \cap B} \mathbb{1}_{\overline{A-k}} d\tilde{\mu} = \overline{\lim}_{n \rightarrow \infty} \frac{1}{|I_n|} \sum_{k \in I_n \cap B} \mu(A - k) = d^*(A) \cdot d^*(B),$$

thus there exists  $p \in \beta\mathbb{Z}$  such that  $d^*(A) \cdot d^*(B) \leq f(p)$ .  $\square$

The above application of Fatou's Lemma is inspired by the proof of [Ber85, Theorem 1.1].

*Proof of Theorem 1.* Assume that  $d^*(A), d^*(B) > 0$ . According to Lemma 2, pick an ultrafilter  $p$  such that  $C := (A - p) \cap B$  has positive upper Banach density. Then  $S := (A - p) - B \supseteq C - C$  is syndetic. Also  $s \in (A - p) - B \implies A - B - s \in p$ . Thus for each finite set  $\{s_1, \dots, s_n\} \subseteq (A - p) - B$ , we have  $\bigcap_{i=1}^n A - B - s_i \in p$ . In particular this intersection is non-empty, hence there exists  $t \in \mathbb{Z}$  such that  $t + \{s_1, \dots, s_n\} \subseteq A - B$ . Summing up, we find that  $A - B$  is piecewise syndetic since it contains shifted copies of all finite subsets of the syndetic set  $(A - p) - B$ .  $\square$

## 2 Jin's Theorem in countable amenable (semi-) groups

Following Jin's original work, it was shown in [JK03] that Theorem 1 is valid in a certain class of abelian groups (including in particular  $\mathbb{Z}^d$ ) and in [Jin08] that it holds in  $\bigoplus_{i=1}^{\infty} \mathbb{Z}$ . Answering a question posed in [JK03] it is proved in [BBF, Theorem 2] that Jin's theorem extends to all countable groups in which the notion of upper Banach density can naturally be formulated, that is, to all countable amenable groups.

There is overwhelming evidence (see in particular [HS98]) that whenever ultrafilters can be used to prove a certain combinatorial statement, then this ultrafilter proof will automatically work in a quite abstract setup. In this spirit the approach of Section 1 effortlessly yields the just mentioned strengthenings of Jin's Theorem; in fact it is not even necessary to restrict the setting to groups.

If a semigroup  $(S, \cdot)$  admits left- and right Følner sequences<sup>1</sup>, the notions  $d_L^*, d_R^*$  of left- resp. right upper Banach density are defined analogously to upper Banach density in  $\mathbb{Z}$ , but with left- resp. right Følner sequences taking the role of intervals  $\{n, \dots, m\}$ . Arguing precisely as in Section 1 we then get that for all  $A, B \subseteq S$  there is an ultrafilter  $p$  on  $S$  such that  $d_L^*(B \cap (q^{-1}A)) = d_L^*(B \cap \{s : As^{-1} \in p\}) \geq d_R^*(A) \cdot d_L^*(B)$ . Consequently we have:

**Theorem 3.** *Let  $(S, \cdot)$  be a semigroup which admits left- and right Følner sequences and let  $A, B \subseteq S$ ,  $d_R^*(A), d_L^*(B) > 0$ . Then  $AB^{-1} = \{s \in S : \exists b \in B \text{ sb} \in A\}$  is (right) piecewise syndetic.*

A subset  $P$  of a semigroup  $(S, \cdot)$  is (right) piecewise syndetic if there exists a finite set  $K \subseteq S$  such that for each finite set  $F \subseteq S$  there is some  $t \in S$  such that  $tF \subseteq PK^{-1}$ .

If  $(S, \cdot)$  is an amenable group, Theorem 3 is equivalent to [BBF, Theorem 2] which asserts that in this setup  $AB$  is (right) piecewise syndetic provided that  $d_R^*(A), d_R^*(B) > 0$ .

## 3 Connections with Bohr sets

So far, our results about the structure of  $A - B$  originated from the fact that for  $d^*(C) > 0$  the set  $C - C$  is syndetic. Følner ([Føl54a, Føl54b]) proved a much stronger assertion, namely that  $C - C$  is "almost" a Bohr<sub>0</sub> set. Bohr<sub>0</sub> sets are the neighborhoods of 0 in the Bohr topology. (Equivalently,  $U \subseteq \mathbb{Z}$  is a Bohr<sub>0</sub> set iff there exist  $\alpha_1, \dots, \alpha_n \in \mathbb{T} = \mathbb{R}/\mathbb{Z}$  and  $\varepsilon > 0$  such that  $\{k \in \mathbb{Z} : \|k\alpha_1\|, \dots, \|k\alpha_n\| < \varepsilon\} \subseteq U$ .) A Bohr set is a translate of a

<sup>1</sup>A sequence  $(F_n)_{n \in \mathbb{N}}$  of finite sets in a semigroup  $S$  is a left/right Følner sequence if  $\lim_{n \rightarrow \infty} |(sF_n) \Delta F_n|/|F_n| = 0$  resp.  $\lim_{n \rightarrow \infty} |(F_n s) \Delta F_n|/|F_n| = 0$  for all  $s \in S$ . The existence of Følner sequences is sometimes used to define if a countable group / semigroup is amenable. We note that all abelian semigroups and all solvable groups fall in this class. See for instance [Pat88].

Bohr<sub>0</sub> set. Every Bohr set is syndetic, but the converse fails badly. Følner showed that if  $d^*(C) > 0$ , then there exist a Bohr<sub>0</sub> set  $U$  and set  $N \subseteq \mathbb{Z}$  with  $d^*(N) = 0$  such that  $C - C \supseteq U \setminus N$ .

In analogy to piecewise syndetic sets Bergelson, Furstenberg and Weiss introduced piecewise Bohr sets. A set  $P$  is piecewise Bohr if it is Bohr on arbitrarily large intervals, that is, if there exist a Bohr set  $U$  and a sequence of intervals  $I_n \subseteq \mathbb{Z}, |I_n| \uparrow \infty$  such that  $P \supseteq U \cap \bigcup_{n=1}^{\infty} I_n$ . Følner's Theorem,  $d^*(C) > 0$  trivially implies that  $C - C$  is piecewise Bohr. We reprove the following refinement of Jin's Theorem obtained in [BFW06].

**Theorem 4.** *Let  $A, B \subseteq \mathbb{Z}, d^*(A), d^*(B) > 0$ . Then  $A - B$  is piecewise Bohr.*

*Proof.* As in the case of piecewise syndetic sets one readily shows that if a set  $P$  contains a translate of every finite subset of a piecewise Bohr set, then  $P$  is piecewise Bohr itself. In Section 1 we have seen that if  $d^*(A), d^*(B) > 0$ , then  $A - B$  contains shifts of all finite pieces of a set  $C - C, d^*(C) > 0$ . By Følner's Theorem the latter set is piecewise Bohr, hence  $A - B$  is piecewise Bohr as well.  $\square$

In the spirit of Section 2 it is possible to proceed along these lines in an abstract countable amenable group. This then leads to the amenable analog of Theorem 4 derived in [BBF, Theorem 3].

It is natural to ask whether one can make assertions about the combinatorial richness of the set  $A - B$  beyond the fact that it is piecewise Bohr. In a certain sense this is not possible. For every piecewise Bohr set  $P \subseteq \mathbb{Z}$ , there exist sets  $A, B \subseteq \mathbb{Z}$  with  $d^*(A), d^*(B) > 0$  such that  $A - B \subseteq P$  ([BBF, Theorem 4]).

## References

- [BBF] Mathias Beiglböck, Vitaly Bergelson, and Alexander Fish. Sumset phenomenon in countable amenable groups. *Adv. Math.*, to appear.
- [Ber85] Vitaly Bergelson. Sets of recurrence of  $\mathbb{Z}^m$ -actions and properties of sets of differences in  $\mathbb{Z}^m$ . *J. London Math. Soc. (2)*, 31(2):295–304, 1985.
- [Ber06] Vitaly Bergelson. Combinatorial and Diophantine applications of ergodic theory. In *Handbook of dynamical systems. Vol. 1B*, pages 745–869. Elsevier B. V., Amsterdam, 2006. Appendix A by A. Leibman and Appendix B by Anthony Quas and Máté Wierdl.
- [BFW06] Vitaly Bergelson, Hillel Furstenberg, and Benjamin Weiss. Piecewise-Bohr sets of integers and combinatorial number theory. In *Topics in discrete mathematics*, volume 26 of *Algorithms Combin.*, pages 13–37. Springer, Berlin, 2006.
- [Føl54a] Erling Følner. Generalization of a theorem of Bogoliouboff to topological abelian groups. With an appendix on Banach mean values in non-abelian groups. *Math. Scand.*, 2:5–18, 1954.

- [Føl54b] Erling Følner. Note on a generalization of a theorem of Bogoliouboff. *Math. Scand.*, 2:224–226, 1954.
- [HS98] Neil Hindman and Dona Strauss. *Algebra in the Stone-Čech compactification*, volume 27 of *de Gruyter Expositions in Mathematics*. Walter de Gruyter & Co., Berlin, 1998. Theory and applications.
- [Jin02] Renling Jin. The sunset phenomenon. *Proc. Amer. Math. Soc.*, 130(3):855–861, 2002.
- [Jin08] Renling Jin. *private communication*, 2008.
- [JK03] Renling Jin and H. Jerome Keisler. Abelian groups with layered tiles and the sunset phenomenon. *Trans. Amer. Math. Soc.*, 355(1):79–97, 2003.
- [Pat88] Alan L. T. Paterson. *Amenability*, volume 29 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1988.