

# Analytic Geometry: Points, Lines, Parabolas and Circles

## Exercise sheet 2 – with step-by-step solutions

Caboara

### 1 Distance between two points

**Reminder.** The distance between  $A(x_1, y_1)$  and  $B(x_2, y_2)$  is

$$d(A, B) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}.$$

**Exercise 1.1.** Compute the distance between  $A(-2, 1)$  and  $B(4, 9)$ .

*Solution. Step 1: differences of coordinates.*  $x_2 - x_1 = 4 - (-2) = 6$ ,  $y_2 - y_1 = 9 - 1 = 8$ .

*Step 2: squares and sum.*  $6^2 + 8^2 = 36 + 64 = 100$ .

*Step 3: square root.*  $d(A, B) = \sqrt{100} = 10$ . □

**Exercise 1.2.** Compute the distance between  $P(3, -4)$  and  $Q(-1, 2)$ .

*Solution. Step 1.*  $x_Q - x_P = -1 - 3 = -4$ ,  $y_Q - y_P = 2 - (-4) = 6$ .

*Step 2.*  $(-4)^2 + 6^2 = 16 + 36 = 52$ .

*Step 3.*  $d(P, Q) = \sqrt{52} = \sqrt{4 \cdot 13} = 2\sqrt{13} \approx 7.21$ . □

### 2 Circle through three points

**Method.** Write the circle in the general form

$$x^2 + y^2 + ax + by + c = 0.$$

Substituting the coordinates of the three points gives a linear system in  $a, b, c$ . If the system has a (unique) solution, the circle has

$$\text{center } C\left(-\frac{a}{2}, -\frac{b}{2}\right), \quad r = \sqrt{\frac{a^2}{4} + \frac{b^2}{4} - c}.$$

If the three points are collinear the system is impossible: no circle passes through them.

**Exercise 2.1.** Find the circle through  $A(1, 5)$ ,  $B(3, 1)$ ,  $D(-6, -2)$ .

*Solution. Step 1: substitute the points.*

$$A(1, 5): 1 + 25 + a + 5b + c = 0 \quad \implies \quad a + 5b + c = -26$$

$$B(3, 1): 9 + 1 + 3a + b + c = 0 \quad \implies \quad 3a + b + c = -10$$

$$D(-6, -2): 36 + 4 - 6a - 2b + c = 0 \quad \implies \quad -6a - 2b + c = -40$$

*Step 2: eliminate  $c$ .*

$$(1\text{st}) - (2\text{nd}): -2a + 4b = -16 \quad \implies \quad a = 2b + 8,$$

$$(2\text{nd}) - (3\text{rd}): 9a + 3b = 30 \quad \implies \quad 3a + b = 10.$$

*Step 3: solve.*  $3(2b + 8) + b = 10 \Rightarrow 7b = -14 \Rightarrow b = -2$ , hence  $a = 4$ . From the second equation:  $c = -10 - 3a - b = -10 - 12 + 2 = -20$ .

Step 4: equation, center, radius.

$$x^2 + y^2 + 4x - 2y - 20 = 0, \quad C(-2, 1), \quad r = \sqrt{4 + 1 + 20} = \sqrt{25} = 5.$$

Check:  $d(C, A) = \sqrt{3^2 + 4^2} = 5$ ,  $d(C, B) = \sqrt{5^2 + 0^2} = 5$ ,  $d(C, D) = \sqrt{(-4)^2 + (-3)^2} = 5$ .  $\square$

**Exercise 2.2.** Find the circle through  $A(5, 2)$ ,  $B(0, 1)$ ,  $D(1, -4)$ .

*Solution.* Step 1: substitute the points.

$$\begin{aligned} A(5, 2) : \quad 25 + 4 + 5a + 2b + c &= 0 & \implies & 5a + 2b + c = -29 \\ B(0, 1) : \quad 0 + 1 + 0 + b + c &= 0 & \implies & b + c = -1 \\ D(1, -4) : \quad 1 + 16 + a - 4b + c &= 0 & \implies & a - 4b + c = -17 \end{aligned}$$

Step 2: eliminate  $c$  by subtracting the second equation from the other two:

$$\begin{aligned} 5a + b &= -28, \\ a - 5b &= -16 & \implies & a = 5b - 16. \end{aligned}$$

Step 3: solve.  $5(5b - 16) + b = -28 \Rightarrow 26b = 52 \Rightarrow b = 2$ , so  $a = -6$ . Then  $c = -1 - b = -3$ .

Step 4: equation, center, radius.

$$x^2 + y^2 - 6x + 2y - 3 = 0, \quad C(3, -1), \quad r = \sqrt{9 + 1 + 3} = \sqrt{13}.$$

Check:  $d(C, A)^2 = 2^2 + 3^2 = 13$ ,  $d(C, B)^2 = (-3)^2 + 2^2 = 13$ ,  $d(C, D)^2 = (-2)^2 + (-3)^2 = 13$ .  $\square$

**Exercise 2.3.** Find, if it exists, the circle through  $A(0, 2)$ ,  $B(1, 5)$ ,  $D(-2, -4)$ .

*Solution.* Step 1: substitute the points.

$$\begin{aligned} A(0, 2) : \quad 4 + 2b + c &= 0 & \implies & 2b + c = -4 \\ B(1, 5) : \quad 26 + a + 5b + c &= 0 & \implies & a + 5b + c = -26 \\ D(-2, -4) : \quad 20 - 2a - 4b + c &= 0 & \implies & -2a - 4b + c = -20 \end{aligned}$$

Step 2: eliminate  $c$  by subtracting the first equation from the other two:

$$\begin{aligned} a + 3b &= -22, \\ -2a - 6b &= -16 & \implies & a + 3b = 8. \end{aligned}$$

Step 3: conclusion. The quantity  $a + 3b$  cannot be both  $-22$  and  $8$ : the system is **impossible**, so **no circle** passes through the three points.

*Geometric reason.* The slopes are  $m_{AB} = \frac{5-2}{1-0} = 3$  and  $m_{AD} = \frac{-4-2}{-2-0} = 3$ : the points are collinear (they lie on the line  $y = 3x + 2$ ), and three distinct collinear points never lie on a circle.  $\square$

### 3 Circle with given center and radius

**Reminder.** The circle with center  $C(x_0, y_0)$  and radius  $r > 0$  is

$$(x - x_0)^2 + (y - y_0)^2 = r^2.$$

**Exercise 3.1.** Write the equation of the circle with center  $C(-4, 1)$  and radius  $r = 3$ .

*Solution.* Step 1: standard form.  $(x + 4)^2 + (y - 1)^2 = 9$ .

Step 2: expand.  $x^2 + 8x + 16 + y^2 - 2y + 1 = 9$ .

Step 3: general form.  $x^2 + y^2 + 8x - 2y + 8 = 0$ .

Check:  $a = 8$ ,  $b = -2$ ,  $c = 8$  give  $C(-4, 1)$  and  $r = \sqrt{16 + 1 - 8} = 3$ .  $\square$

**Exercise 3.2.** Write the equation of the circle with center  $C(\frac{2}{3}, -1)$  and radius  $r = 2$ .

Solution. Step 1: standard form.  $(x - \frac{2}{3})^2 + (y + 1)^2 = 4$ .

Step 2: expand.  $x^2 - \frac{4}{3}x + \frac{4}{9} + y^2 + 2y + 1 = 4$ .

Step 3: general form. Since  $\frac{4}{9} + 1 - 4 = \frac{4-27}{9} = -\frac{23}{9}$ ,

$$x^2 + y^2 - \frac{4}{3}x + 2y - \frac{23}{9} = 0 \quad \text{or, multiplying by 9,} \quad 9x^2 + 9y^2 - 12x + 18y - 23 = 0.$$

Check:  $a = -\frac{4}{3}$ ,  $b = 2$ ,  $c = -\frac{23}{9}$  give  $C(\frac{2}{3}, -1)$  and  $r^2 = \frac{4}{9} + 1 + \frac{23}{9} = \frac{36}{9} = 4$ .  $\square$

## 4 Center and radius from the equation

**Method.** An equation  $\alpha x^2 + \alpha y^2 + \dots = 0$  (same coefficient of  $x^2$  and  $y^2$ , no  $xy$  term) is first divided by  $\alpha$  to get  $x^2 + y^2 + ax + by + c = 0$ . Completing the squares,

$$(x + \frac{a}{2})^2 + (y + \frac{b}{2})^2 = \frac{a^2}{4} + \frac{b^2}{4} - c.$$

It is a circle only if the right-hand side is *positive*.

**Exercise 4.1.** Find center and radius of  $x^2 + y^2 + 6x - 4y - 12 = 0$ .

Solution. Step 1: group.  $(x^2 + 6x) + (y^2 - 4y) - 12 = 0$ .

Step 2: complete the squares.  $x^2 + 6x = (x + 3)^2 - 9$ ,  $y^2 - 4y = (y - 2)^2 - 4$ , so

$$(x + 3)^2 - 9 + (y - 2)^2 - 4 - 12 = 0 \implies (x + 3)^2 + (y - 2)^2 = 25.$$

Step 3: read off. Since  $25 > 0$ : center  $C(-3, 2)$ , radius  $r = 5$ .

With the formulas:  $a = 6$ ,  $b = -4$ ,  $c = -12$ :  $C(-3, 2)$ ,  $r^2 = 9 + 4 + 12 = 25$ .  $\square$

**Exercise 4.2.** Find center and radius of  $4x^2 + 4y^2 - 4x + 12y - 6 = 0$ .

Solution. Step 1: normalize. The coefficients of  $x^2$  and  $y^2$  are equal; divide by 4:

$$x^2 + y^2 - x + 3y - \frac{3}{2} = 0.$$

Step 2: complete the squares.  $x^2 - x = (x - \frac{1}{2})^2 - \frac{1}{4}$ ,  $y^2 + 3y = (y + \frac{3}{2})^2 - \frac{9}{4}$ , so

$$(x - \frac{1}{2})^2 + (y + \frac{3}{2})^2 = \frac{1}{4} + \frac{9}{4} + \frac{3}{2} = \frac{16}{4} = 4.$$

Step 3: read off. Center  $C(\frac{1}{2}, -\frac{3}{2})$ , radius  $r = 2$ .

Warning: applying the formulas to the *non-divided* equation would give a wrong result.  $\square$

**Exercise 4.3.** Find, if possible, center and radius of  $x^2 + y^2 + 4x - 6y + 15 = 0$ .

Solution. Step 1: complete the squares.  $x^2 + 4x = (x + 2)^2 - 4$ ,  $y^2 - 6y = (y - 3)^2 - 9$ , so

$$(x + 2)^2 - 4 + (y - 3)^2 - 9 + 15 = 0 \implies (x + 2)^2 + (y - 3)^2 = -2.$$

Step 2: conclusion. The left-hand side is a sum of squares, hence  $\geq 0$  for every real  $(x, y)$ , while the right-hand side is  $-2 < 0$ . No real point satisfies the equation: it is **not a circle**.

With the formula:  $r^2 = \frac{a^2}{4} + \frac{b^2}{4} - c = 4 + 9 - 15 = -2 < 0$ .

Remark. With constant term 13 instead of 15 we would get  $(x + 2)^2 + (y - 3)^2 = 0$ , satisfied only by the point  $(-2, 3)$ : a “circle of radius 0”, again not a genuine circle.  $\square$

## 5 Intersections of a line and a circle


**Method.** Solve the system circle–line by substitution: one obtains a quadratic equation, whose discriminant  $\Delta$  tells us the number of intersection points:

$\Delta > 0$  : 2 points (secant),       $\Delta = 0$  : 1 point (tangent),       $\Delta < 0$  : no points (external line).

Equivalently, compare the distance  $d$  of the center from the line with the radius  $r$ :  $d < r$ ,  $d = r$ ,  $d > r$  respectively. Recall  $d(P_0, \alpha x + \beta y + \gamma = 0) = \frac{|\alpha x_0 + \beta y_0 + \gamma|}{\sqrt{\alpha^2 + \beta^2}}$ .

In all three exercises the circle is

$$\Gamma : x^2 + y^2 + 2x - 6y - 15 = 0, \quad \text{center } C(-1, 3), \quad r = \sqrt{1 + 9 + 15} = 5.$$



**Exercise 5.1.** Find the intersections of  $\Gamma$  with the line  $s : 2x + y - 6 = 0$ .

*Solution. Step 1: solve the line for  $y$ .  $y = 6 - 2x$ .*

*Step 2: substitute into  $\Gamma$ .*

$$\begin{aligned} x^2 + (6 - 2x)^2 + 2x - 6(6 - 2x) - 15 &= 0 \\ \implies x^2 + 36 - 24x + 4x^2 + 2x - 36 + 12x - 15 &= 0 \\ \implies 5x^2 - 10x - 15 &= 0. \end{aligned}$$

*Step 3: solve.* Dividing by 5:  $x^2 - 2x - 3 = 0$ , with  $\Delta = 4 + 12 = 16 > 0$ :

$$x = \frac{2 \pm 4}{2} \implies x_1 = 3, \quad x_2 = -1.$$

*Step 4: find  $y$ .*  $y_1 = 6 - 6 = 0$ ,  $y_2 = 6 + 2 = 8$ . **Two intersection points:**  $(3, 0)$  and  $(-1, 8)$ ; the line is secant.

$$\text{Check with the distance: } d(C, s) = \frac{|-2 + 3 - 6|}{\sqrt{4 + 1}} = \frac{5}{\sqrt{5}} = \sqrt{5} \approx 2.24 < 5. \quad \square$$

**Exercise 5.2.** Find the intersections of  $\Gamma$  with the line  $t : 3x - 4y - 10 = 0$ .

*Solution. Step 1: solve the line for  $y$ .  $y = \frac{3x - 10}{4}$ .*

*Step 2: substitute into  $\Gamma$  and multiply everything by 16 to clear denominators:*

$$\begin{aligned} 16x^2 + (3x - 10)^2 + 32x - 24(3x - 10) - 240 &= 0 \\ \implies 16x^2 + 9x^2 - 60x + 100 + 32x - 72x + 240 - 240 &= 0 \\ \implies 25x^2 - 100x + 100 &= 0. \end{aligned}$$

*Step 3: solve.* Dividing by 25:  $x^2 - 4x + 4 = 0$ , i.e.  $(x - 2)^2 = 0$ , so  $\Delta = 0$  and  $x = 2$  (double root).

*Step 4: find y.*  $y = \frac{6 - 10}{4} = -1$ . **One intersection point:**  $(2, -1)$ ; the line is tangent to  $\Gamma$ .

*Check with the distance:*  $d(C, t) = \frac{|-3 - 12 - 10|}{\sqrt{9 + 16}} = \frac{25}{5} = 5 = r$ .  $\square$

**Exercise 5.3.** Find the intersections of  $\Gamma$  with the line  $e : x - y + 12 = 0$ .

*Solution. Step 1: solve the line for y.*  $y = x + 12$ .

*Step 2: substitute into  $\Gamma$ .*

$$\begin{aligned} x^2 + (x + 12)^2 + 2x - 6(x + 12) - 15 &= 0 \\ \implies x^2 + x^2 + 24x + 144 + 2x - 6x - 72 - 15 &= 0 \\ \implies 2x^2 + 20x + 57 &= 0. \end{aligned}$$

*Step 3: discriminant.*  $\Delta = 20^2 - 4 \cdot 2 \cdot 57 = 400 - 456 = -56 < 0$ .

*Conclusion.* The equation has no real solutions: **no intersection points**; the line is external to  $\Gamma$ .

*Check with the distance:*  $d(C, e) = \frac{|-1 - 3 + 12|}{\sqrt{2}} = \frac{8}{\sqrt{2}} = 4\sqrt{2} \approx 5.66 > 5$ .  $\square$

## 6 Intersections of a parabola and a circle


**Method.** In this sheet the parabolas open *downwards*:  $y = q - x^2$ . Replace  $x^2$  with  $q - y$  in the equation of the circle (center on the  $y$ -axis). We get a quadratic equation in  $y$  only. Then, for each real root  $\bar{y}$ :

- if  $\bar{y} < q$ :  $x^2 = q - \bar{y} > 0$  gives *two* points  $(\pm\sqrt{q - \bar{y}}, \bar{y})$ ;
- if  $\bar{y} = q$ :  $x^2 = 0$  gives *one* point  $(0, q)$  (the vertex);
- if  $\bar{y} > q$ :  $x^2 = q - \bar{y} < 0$  gives *no* points.

Note that the condition is now  $\bar{y} \leq q$  (the parabola lies *below* its vertex).

**Non-symmetric configurations (Exercises 6.3–6.5).** When the axis of the parabola does not pass through the center of the circle, the trick above does not work. Substitute instead the whole expression  $y = -x^2 + bx + c$  into the equation of the circle: we get a polynomial equation of degree 4 in  $x$ .

- Look for integer roots among the divisors of the constant term, and lower the degree by polynomial long division (or by factoring out  $x$ , or by grouping).
- Each *real* root  $\bar{x}$  gives exactly *one* point  $(\bar{x}, -\bar{x}^2 + b\bar{x} + c)$ .
- A *double* root corresponds to a point where the two curves are tangent.



**Exercise 6.1.** Find the intersections of the circle  $x^2 + y^2 = 25$  and the parabola  $y = -x^2 - 7$ .

*Solution. Step 1: express  $x^2$  from the parabola.  $x^2 = -7 - y$ , with the condition  $y \leq -7$ .*

*Step 2: substitute into the circle.*

$$(-7 - y) + y^2 = 25 \implies y^2 - y - 32 = 0.$$

*Step 3: solve.  $\Delta = 1 + 128 = 129 > 0$ , so there are two real roots*

$$y_{1,2} = \frac{1 \pm \sqrt{129}}{2}, \quad y_1 \approx 6.18, \quad y_2 \approx -5.18.$$

*Step 4: check the condition  $y \leq -7$ .  $y_1 > 0 > -7$  is clearly rejected. For  $y_2$ :*

$$\frac{1 - \sqrt{129}}{2} \leq -7 \iff 1 - \sqrt{129} \leq -14 \iff \sqrt{129} \geq 15 \iff 129 \geq 225,$$

which is false, so  $y_2 > -7$  is rejected too.

*Conclusion. No intersection points*, even though  $\Delta > 0$ : both roots give  $x^2 = -7 - y < 0$ .

*Quick check.* Every point of the parabola has  $y = -x^2 - 7 \leq -7$ , while every point of the circle has  $|y| \leq 5$ : the two curves cannot meet.  $\square$

**Exercise 6.2.** Find the intersections of the circle  $x^2 + y^2 = 25$  and the parabola  $y = -x^2 - 5$ .

*Solution. Step 1.  $x^2 = -5 - y$ , with the condition  $y \leq -5$ .*

*Step 2: substitute.*

$$(-5 - y) + y^2 = 25 \implies y^2 - y - 30 = 0.$$

*Step 3: solve.  $\Delta = 1 + 120 = 121$ ,  $y = \frac{1 \pm 11}{2}$ , so  $y_1 = 6$ ,  $y_2 = -5$ .*

*Step 4: check the condition  $y \leq -5$  and find  $x$ .*

- $y_1 = 6 > -5$ :  $x^2 = -11$ , impossible: no points;

- $y_2 = -5$ :  $x^2 = 0 \Rightarrow x = 0$ : the point  $(0, -5)$ .

*Conclusion.* **One intersection point:**  $(0, -5)$ . The vertex of the parabola is the lowest point of the circle; the parabola opens downwards, away from the circle, and the two curves are tangent there.  $\square$

**Exercise 6.3.** Find the intersections of the circle  $(x + 1)^2 + y^2 = 10$  and the parabola  $y = -x^2 + 3x - 3$ .

*Solution.* *Step 1: substitute  $y$  into the circle.* We need  $(-x^2 + 3x - 3)^2 = x^4 - 6x^3 + 15x^2 - 18x + 9$  and  $(x + 1)^2 = x^2 + 2x + 1$ , so

$$\begin{aligned} x^2 + 2x + 1 + x^4 - 6x^3 + 15x^2 - 18x + 9 &= 10 \\ \implies x^4 - 6x^3 + 16x^2 - 16x &= 0. \end{aligned}$$

*Step 2: factor out  $x$ .*  $x(x^3 - 6x^2 + 16x - 16) = 0$ , so  $x = 0$  is a root.

*Step 3: roots of the cubic.* Integer roots must divide 16: try  $\pm 1, \pm 2, \dots$ . For  $x = 1$ :  $1 - 6 + 16 - 16 = -5 \neq 0$ ; for  $x = 2$ :  $8 - 24 + 32 - 16 = 0$ . Divide by  $x - 2$ :

$$\begin{array}{r|l} x^3 - 6x^2 + 16x - 16 & x - 2 \\ -x^3 + 2x^2 & \hline -4x^2 + 16x & \\ 4x^2 - 8x & \\ \hline 8x - 16 & \\ -8x + 16 & \\ \hline 0 & \end{array}$$

so  $x^3 - 6x^2 + 16x - 16 = (x - 2)(x^2 - 4x + 8)$ .

*Step 4: the remaining quadratic.*  $x^2 - 4x + 8 = 0$  has  $\Delta = 16 - 32 = -16 < 0$ : no real roots.

*Step 5: find  $y$ .* The real roots are  $x = 0$  and  $x = 2$ :

$$x = 0 : y = -3, \quad x = 2 : y = -4 + 6 - 3 = -1.$$

*Conclusion.* **Two intersection points:**  $(0, -3)$  and  $(2, -1)$ .

*Check:*  $(0 + 1)^2 + (-3)^2 = 10$  and  $(2 + 1)^2 + (-1)^2 = 10$ .  $\square$

**Exercise 6.4.** Find the intersections of the circle  $x^2 + y^2 = 5$  and the parabola  $y = -x^2 + 2x + 1$ .

*Solution.* *Step 1: substitute  $y$  into the circle.* Since  $(-x^2 + 2x + 1)^2 = x^4 - 4x^3 + 2x^2 + 4x + 1$ ,

$$\begin{aligned} x^2 + x^4 - 4x^3 + 2x^2 + 4x + 1 &= 5 \\ \implies x^4 - 4x^3 + 3x^2 + 4x - 4 &= 0. \end{aligned}$$

*Step 2: a first root.* Integer roots must divide 4: candidates  $\pm 1, \pm 2, \pm 4$ . For  $x = 1$ :  $1 - 4 + 3 + 4 - 4 = 0$ . Divide by  $x - 1$  (note the missing  $x$  term in the quotient):

$$\begin{array}{r|l} x^4 - 4x^3 + 3x^2 + 4x - 4 & x - 1 \\ -x^4 + x^3 & \hline -3x^3 + 3x^2 & \\ 3x^3 - 3x^2 & \\ \hline 4x - 4 & \\ -4x + 4 & \\ \hline 0 & \end{array}$$

so the equation becomes  $(x - 1)(x^3 - 3x^2 + 4) = 0$ .

*Step 3: a second root.* For  $x = -1$ :  $-1 - 3 + 4 = 0$ . Divide by  $x + 1$ :

$$\begin{array}{r|l}
 x^3 - 3x^2 & + 4 \\
 -x^3 & -x^2 \\
 \hline
 & -4x^2 \\
 & 4x^2 + 4x \\
 \hline
 & 4x + 4 \\
 & -4x - 4 \\
 \hline
 & 0
 \end{array}$$

and  $x^2 - 4x + 4 = (x - 2)^2$ . Hence

$$x^4 - 4x^3 + 3x^2 + 4x - 4 = (x - 1)(x + 1)(x - 2)^2.$$

*Step 4: find  $y$ .*

$$x = 1 : y = 2, \quad x = -1 : y = -1 - 2 + 1 = -2, \quad x = 2 : y = -4 + 4 + 1 = 1.$$

*Conclusion. Three intersection points:*  $(1, 2)$ ,  $(-1, -2)$ ,  $(2, 1)$ . The root  $x = 2$  is *double*: at  $(2, 1)$  the parabola and the circle are tangent. Unlike the symmetric case, the tangency point is not the vertex of the parabola (the vertex is  $(1, 2)$ , where the curves simply cross).

*Check:*  $1 + 4 = 5$ ,  $1 + 4 = 5$ ,  $4 + 1 = 5$ . □

**Exercise 6.5.** Find the intersections of the circle  $(x + 2)^2 + y^2 = 25$  and the parabola  $y = -x^2 + 2x + 3$ .

*Solution. Step 1: substitute  $y$  into the circle.* Since  $(-x^2 + 2x + 3)^2 = x^4 - 4x^3 - 2x^2 + 12x + 9$  and  $(x + 2)^2 = x^2 + 4x + 4$ ,

$$\begin{aligned}
 x^2 + 4x + 4 + x^4 - 4x^3 - 2x^2 + 12x + 9 &= 25 \\
 \implies x^4 - 4x^3 - x^2 + 16x - 12 &= 0.
 \end{aligned}$$

*Step 2: a first root.* Integer roots must divide 12. For  $x = 1$ :  $1 - 4 - 1 + 16 - 12 = 0$ . Divide by  $x - 1$ :

$$\begin{array}{r|l}
 x^4 - 4x^3 & -x^2 + 16x - 12 \\
 -x^4 & +x^3 \\
 \hline
 & -3x^3 - x^2 \\
 & 3x^3 - 3x^2 \\
 \hline
 & -4x^2 + 16x \\
 & 4x^2 - 4x \\
 \hline
 & 12x - 12 \\
 & -12x + 12 \\
 \hline
 & 0
 \end{array}$$

so the equation becomes  $(x - 1)(x^3 - 3x^2 - 4x + 12) = 0$ .

*Step 3: factor the cubic by grouping.*

$$x^3 - 3x^2 - 4x + 12 = x^2(x - 3) - 4(x - 3) = (x - 3)(x^2 - 4) = (x - 3)(x - 2)(x + 2).$$

Hence the four roots are  $x = 1, 2, 3, -2$ .

*Step 4: find  $y$ .*

$$x = 1 : y = 4, \quad x = 2 : y = 3, \quad x = 3 : y = 0, \quad x = -2 : y = -4 - 4 + 3 = -5.$$

*Conclusion. Four intersection points:*  $(1, 4)$ ,  $(2, 3)$ ,  $(3, 0)$ ,  $(-2, -5)$ . Note that, without symmetry, the points no longer come in pairs  $(\pm \bar{x}, \bar{y})$ .

*Check:*  $9 + 16$ ,  $16 + 9$ ,  $25 + 0$ ,  $0 + 25$  all equal 25. □