


# Hitchin fibrations and related combinatorics


## PLAN

- Moduli space  $M_{n,d}$
- Hitchin fibration  $\mathcal{X} \downarrow$
- $\mathcal{X}^{-1}(a)$  BNR correspondence  $A_n$
- Intersection Cohomology  $M_{n,d}$

SETTING Let  $C$  be a smooth proj alg curve/c  
(a Riemann surface)



DEF a vector bundle  $E$  over  $C$  of rank  $n$



Operations on vector bundles:

- $\oplus$  direct sum,  $\otimes$  tensor product, dual,  $\wedge^n$  exterior power
- $S^\lambda$  Schur functors


EG  $\Lambda^n E$  is a line bundle  $m=1$

$\stackrel{!!}{\text{det}} E$

DEF Let  $L$  be a line bundle  $s$  a meromorphic section



$\text{deg } L := \# \text{ zeros } (s) - \# \text{ poles } (s)$



EX  $\text{deg}(L)$  does not depend on the choice of  $s$

DEF  $\text{deg}(E) := \text{deg}(\text{det } E)$

(  $\text{deg } E = \int_C c_1(E)$  first Chern class )

DEF Tangent bundle to  $X (= C)$


$T^*X$  be the cotangent bundle

$(T^*X)^\vee$   $\omega_X := \text{det}(T^*X)$  is called canonical

DEF A Higgs bundle is a pair  $(E, \phi)$  where

$\phi: E \rightarrow E \otimes \omega_C$





$\phi_p \in \text{Mat}_{n,n}(\mathbb{C})$

$\phi|_U$  has entries that depends on  $U \cong \mathbb{C} = \mathbb{A}^1$

$U \cong \mathbb{C} = \mathbb{A}^1$

$\phi|_U \in \text{Mat}_{n \times n}(\mathbb{C}[t])$

we glue together the local patches the transition functions behaves "like"  $w_C$

GOAL DEFINE The moduli space of Higgs bundles

EX 1  $GL_n \curvearrowright \text{Mat}_{n \times n}$  by conjugation

$\text{Mat}_{n,n} / GL_n = \{ \text{Jordan normal forms} \}$

endow with quotient topology

$\begin{pmatrix} \lambda & & & \\ & \lambda & & \\ & & \lambda & \\ & & & \lambda \end{pmatrix} \approx J_{\lambda, n}$  for  $t \neq 0$   
 $n_1 + n_2 = n$

$t \rightarrow 0 \quad J_{\lambda, n_1} \oplus J_{\lambda, n_2}$

$GL_n \curvearrowright J_{\lambda, n} \ni J_{\lambda, n_1} \oplus J_{\lambda, n_2}$

$\Rightarrow \text{Mat}_{n,n} / GL_n$  is not separated (not  $T_2$ )

$A \in \text{Mat}_{n,n}$  define a closed point  $[A]$

iff  $A$  is diagonalizable

$$j_{\lambda, n} \cong j_{\lambda, n_1} \oplus j_{\lambda, n_2} \oplus \dots$$


$$0 \rightarrow W \rightarrow V \rightarrow V/W \rightarrow 0$$

$\downarrow$   $\downarrow$   $\downarrow$   
 $j_{\lambda, n_1}$   $j_{\lambda, n_2}$   $j_{\lambda, n_2}$

impose  $V \cong W \oplus V/W$

Mat<sub>n,n</sub> Gln is an alg var.


Example 2  $C = C_1 \cup C_2$



{ rank 2 torsion free sheaves of degree d }

$$\begin{array}{c} \downarrow \\ \mathbb{Z} \end{array} \quad \begin{array}{c} F \\ \downarrow \\ \deg(F|_{C_1}) \end{array}$$

$$\deg(F|_{C_2}) = d - \deg(F|_{C_1})$$



infinitely many irreducible components

is not of finite type

solution Impose a stability condition

eg  $\deg(F|_{C_1}) \leq d$

$$\deg(F|_{C_2}) \leq d.$$

DEF  $(E, \phi)$  Higgs bundle

$(F, \phi|_F)$  is a sub Higgs-bundle if

$F \subset E$  is a sub bundle

and  $F$  is  $\phi$  invariant

$$\mu(E) := \frac{\deg(E)}{\text{rank}(E)} = \frac{d}{n} \in \mathbb{Q} \quad \text{slope of } E$$

DEF  $(E, \phi)$  is (semi)stable Higgs bundle if

$\forall F \subset E$  sub Higgs-bundle

$$\mu(F) < \mu(E) \quad (\leq)$$

DEF  $M_{n,d}^c := \{ \text{semi stable Higgs bundle of rank } n \text{ and degree } d \}$

S-equivalence

$$0 \rightarrow F \rightarrow E \rightarrow G \rightarrow 0 \quad \begin{array}{l} \text{semi stable} \\ \text{s.e.s. of } \bigvee \text{ Higgs bundles} \end{array}$$

$$\mu(F) = \mu(E) = \mu(G)$$

$$\Rightarrow F \oplus G \cong E \quad \begin{array}{l} \text{S-equivalent} \\ \text{sub-bundle} \end{array}$$

EX  $\mu(F) = \mu(E) \quad F \subset E \quad E \text{ s.s.} \Rightarrow$

$F \text{ is s.s.} \quad \mu(E/F) = \mu(E) \quad E/F \text{ is s.s.}$

REMARK .  $\exists (E, \phi)$  semi stable s.t.  $E$  is

not stable as vector bundle

- Rank, degree are additive on s.e.s.
- $N_{d,n}^C$  moduli space of vector bundles
- $E$  a s.s. vector bundle

$$T_E^x N_{d,n}^C \cong H^2(C, \text{End}(E))^V \underset{\text{Serre duality}}{\cong} H^0(C, \text{End}(E) \otimes \omega_C)$$

$\Downarrow$   
 $\phi: E \rightarrow E \otimes \omega_C$

$$T^x N_{d,n}^C \subseteq M_{d,n} \quad \text{open set}$$

$$(E, \phi) \quad (E, \phi)$$

$E$  is s.s. vector bundle

$\Rightarrow \forall \phi \quad (E, \phi)$  is s.s. Higgs bundle

$$\dim M_{d,n} = 2 \dim N_{d,n}^C = 2n^2(g-1) + 2$$


•  $M_{d,n}$  is non-compact, singular

• if  $(d,n) = 1$  then  $M_{d,n}$  is smooth

Ex  $(d,n) = 1$   $(E, \phi)$  s.s. <sup>Higgs</sup>  $\Rightarrow (E, \phi)$  is stable <sup>Higgs</sup>

Hitchin fibration

$(E, \phi) \rightsquigarrow \chi_\phi$  characteristic polynomial



$$\mathcal{X}_\phi(\lambda) \in \mathbb{C}[\lambda] \quad \mathcal{X}_{\phi|_U}(\lambda) \in \underbrace{\mathbb{C}[t][\lambda]}_{O(U)}$$

"The entries of  $\phi$  are sections of  $w_C$ "

$$\text{Tr}(\phi) \in H^0(C, w_C) \quad \det \phi \in H^0(C, w_C^{\otimes n})$$

$$\mathcal{X}_\phi(\lambda) = \lambda^n + s_1 \lambda^{n-1} + \dots + s_n$$

$$s_i \in H^0(C, w_C^{\otimes i})$$

DEF

$$A_n := \bigoplus_{i=1}^n H^0(C, w_C^{\otimes i}) \cong \mathbb{A}^n$$

EX USE R-R. to prove  $N = n^2(g-1) + 1$

The Hitchin fibration ~~map~~ map

$$\begin{aligned} \mathcal{X} : M_{n,d} &\longrightarrow A_n \\ (E, \phi) &\longmapsto (\text{coeff } \mathcal{X}_E) \end{aligned}$$

REMARK :  $M_{n,d}$  is not projective,  $\mathcal{X}$  is surjective

•  $\mathcal{X}$  is a proper map




Goal

$$\mathcal{X}^{-1}(a) = ? = \{ (E, \phi) \mid \mathcal{X}_\phi = q_a \}$$

$$a \in A_n \quad q_a(\lambda) = \lambda^n + a_1 \lambda^{n-1} + \dots + a_n$$

$q_a$  can be evaluated at points of  $T^*C$





$$q_a|_p = \lambda^n + (a_2)_p \lambda^{n-1} + \dots + (a_n)_p \in \mathbb{C}$$


$$q_a|_U = \lambda^n + a_2(t) \lambda^{n-1} + \dots + a_n(t)$$

$$q_a = \lambda^n + a_1 \lambda^{n-1} + \dots + a_n$$

$\uparrow$   $s \in H^0(C, \omega_C)$

DEF  $C_a \subset T^*C$

$C_a =$  zero locus of  $q_a(\lambda)$   
 $\uparrow$  very singular




$C_a$  is called The spectral curve


EX  $q_a = P_1^{k_1} \cdot P_2^{k_2} \cdot \dots \cdot P_r^{k_r}$

local

1-  $\lambda^2 + 2t\lambda + t^2 = (\lambda+t)^2$



2-  $\lambda^2 + 3t\lambda + 2t^2 = (\lambda+t)(\lambda+2t)$






$$3 - x^2 - t^3$$

irreducible poly (curve)


singular



4 - generic case

$$x^2 + t^2 - 1$$

smooth and irreducible



$$\mathcal{X}^{-1}(a)$$


---

$$M_{n,d} = \{ (E, \phi) \mid \begin{matrix} \text{rk } n \\ \text{deg } d \end{matrix} \}$$

$$\mathcal{X} \downarrow \\ A_n \ni a$$

$$C_a \subset T^*C \\ \downarrow \text{n-cover} \\ C$$

EX  $C_a$  is smooth

$$g(C_a) = n^2(g-1) + 1$$

Hint 1: adj formula & RR Thm

Hint 2: R-H Thm & resultant

consider a s.t.  $C_a$  is smooth

$L$  line bundle on  $C_a$

$$L \rightarrow C_a$$

$$\downarrow \pi$$

$$\pi_* L \rightarrow C$$

$\leftarrow$  is a vector bundle


eigenspaces  $\phi$

$$\mathbb{C} \rightarrow \bullet$$

$$\mathbb{C} \rightarrow \bullet$$

$$\mathbb{C} \rightarrow \bullet$$

$$\mathbb{C}^n \rightarrow \rho \quad \begin{matrix} \lambda_1, \dots, \lambda_n \\ \text{eigenvalues } \phi \end{matrix}$$



$$\pi^* U$$

$$\downarrow$$



FACT  $\cdot \deg(\pi_* L) = \deg(L) - n(n-1)(g-1)$

$$\leftarrow \det \pi_* L = \text{Norm}(L) \otimes \det \pi_* \mathcal{O}$$

• There is a natural endomorphism on  $\pi_* L$  is given by multiplication  $\lambda$  canonical section

$$\phi_P = \begin{pmatrix} \lambda_1 & & 0 \\ & \lambda_2 & \\ 0 & & \ddots \\ & & & \lambda_n \end{pmatrix}$$

THM Beauville Narasimhan Ramanan

if  $C_a$  is smooth then

$$\mathcal{X}^{-1}(a) \cong \int_{C_a}^{d + n(n-1)(g-1)} = \left\{ \begin{array}{l} \text{Line bundles on } C_a \\ \text{of degree } d + n(n-1)(g-1) \end{array} \right\}$$

$$(\pi_* L, \phi_L) \longleftarrow L$$

$$(E, \phi) \longmapsto ((p, \lambda) \mapsto \text{gen eigenspace of } \phi_p \text{ associate } \lambda)$$

COR  $\dim \mathcal{M}_{n,d} = \dim A_n + \dim J_{C_a} = 2n^2(g-1) + 2$

Stratification of  $A_n$


DEF  $A_n^{\text{red}} \subset A_n$   $A_n^{\text{red}} = \{ \underline{a} \mid q_{\underline{a}} = p_2 \cdots p_r \mid \substack{r \geq 1 \\ k_i \geq 1} \}$   
 $= \{ \underline{a} \mid C_{\underline{a}} \text{ is reduced} \}$

$M_{\text{red}} := \mathcal{X}^{-1}(A_n^{\text{red}})$

$S_{\underline{n}} := \{ \underline{a} \mid q_{\underline{a}} = p_2 \cdots p_r \text{ s.t. } \deg p_i = n_i \}$


$\underline{n} \vdash n := \{ \underline{a} \mid C_{\underline{a}} \text{ has } r \text{ irr. components of degree } n_i \}$   
 $\parallel$   
 $\{ n_1, n_2, \dots, n_r \}$

EG  $n=3$



$q_{\underline{a}} = p_1 p_2 p_3 = t_2 t_2$   
 (with arrows indicating the mapping from p's to t's and a note 'deg 2')

branching around  $S_{2,1,1} \leftrightarrow$  ref partition of  $[3]$




$A_{n_1} \times A_{n_2} \times \dots \times A_{n_k} \rightarrow \overline{S}_{\underline{n}}$   
 $A_1 \times A_1 \times A_1 \rightarrow S_{2,1,1}$   
 $6 \mapsto 1$

Dual graph of  $C_{\underline{a}}$

$\underline{a} \in S_{\underline{n}}$   $\Gamma_{\underline{n}}$  on  $r$  vertices and  $x_{ij}$  edges  $i-j$   
 $x_{ij} := \# C_i \cap C_j = n_i n_j (2g-2) \in \mathbb{N}_+$

$\{ \text{rank 1 torsion free sheaves on } C_{\underline{a}} \}$

$L^{\psi}$



DEF  $L$  is  $\underline{n}$ -stable if  $\chi(L|_{C'}) > \chi(L) \frac{\deg(C' \rightarrow C)}{n} = n_i \quad \forall C' \not\subseteq C_a$

$$\chi(L) = \dim H^0(L) - \dim H^2(L) = \deg(L) - \rho_a + 1$$

PROP  $x_i := \chi(L|_{C_i}) - (1-g)n_i$

$L$  is  $\underline{n}$ -stable iff  $\underline{n} = \{n_1, \dots, n_r\} \vdash n$

$$\sum_{i \in K} x_i > \sum_{i \in K} y_{ij} + \sum_{i \in K} \frac{d n_i}{n} \quad \forall K \subset [r]$$

THM  $\underline{a} \in S_{\underline{n}} \subseteq A_n^{\text{red}}$

$$\mathcal{J}_{\underline{n}\text{-stable}}(C_a) = \left\{ \begin{array}{l} \text{rank 1 t.f. sheaves} \\ \underline{n}\text{-stable of degree } d \end{array} \right\} \xrightarrow{\sim} \mathcal{X}^{-1}(\underline{a})$$

$L \longmapsto (\pi_* L, \chi \cdot)$   
 $\uparrow$   $\leftarrow$   $n$ -stability  $\longleftrightarrow$   $\mu$ -stability  
 has pure dimension  $d(C_a) = n^2(g-1) + 1$

$$\sum_{i \in [n]} x_i = \sum_{i,j \in [n]} y_{ij} + d$$

Q # irr. components of  $\mathcal{J}(C_a) = ?$

are determined by  $(x_1, \dots, x_r)$

$$Z_{n,d} = \left\{ (x_1, \dots, x_r) \in \mathbb{R}^r \mid \begin{array}{l} \sum x_i = \sum y_{ij} + d \\ \sum_{i \in K} x_i > \sum_{i \in K} y_{ij} + \sum_{i \in K} \frac{d n_i}{n} \end{array} \right\}$$

$$\# \text{ irr. components} = \# (Z_{n,d} \cap Z^r) =: C(Z_{n,d})$$

Cor # irr. cap  $\mathcal{X}^{-1}(\underline{a}) = C(Z_{n,d})$

# Combinatorics

Let  $\Gamma$  a graph on  $[r]$





Let  $T \vdash [r]$   $T = \{T_1, T_2, \dots, T_k\}$

deleted graph  $\Gamma_T = \{e \in \Gamma \mid e = i-j \text{ for } k \text{ s.t. } i, j \in T_k\}$

contracted graph  $\Gamma^T = \frac{\Gamma}{\Gamma_T} \cong \{e \in \Gamma \setminus \Gamma_T\}$  on vertex set  $[k]$

eg  $T = 12 | 3$






DEF a flat of  $\Gamma$  is  $T \vdash [r] \leq T$ .


every block  $T_i$  induces a connected subgraph

The flats are ordered by refinement

eg  $T = 1 | 23$



The flats are



## Graphical Zonotopes


$\Gamma$  be a graph

$$Z_\Gamma := \sum_{i,j} x_{ij} [e_i, e_j] \in \mathbb{R}^E$$

Minkowski sum



# edges between  $i$  and  $j$

segment between  $e_i$  and  $e_j$



FACT •  $Z_P = \left\{ (x_1, \dots, x_r) \mid \sum x_i = \sum y_{ij}, \forall k \in [r] \sum_{i \in K} x_i \geq \sum_{i \in K} y_{ij} \right\}$

•  $\{ \text{faces of } Z_P \} \leftrightarrow \left\{ (T, a) \mid \begin{array}{l} T \text{ is a flat of } P \\ a \text{ is an acyclic orientation} \\ \text{of } P^T \end{array} \right\}$



Remark # of rays  $\chi^1(a) = C(Z_{P_n} + w)$

$P_n$  is the dual graph of  $C_n$   $w = (w_i)$   $w_i = \frac{dn_i}{n}$

### Ehrhart Theory

DEF  $\text{ehr}_P(t) := \# (tP \cap \mathbb{Z}^r)$   $t \in \mathbb{N}_+$

eg  $P = [0, \frac{1}{2}] \in \mathbb{R}$

$\text{ehr}_P(t) = \begin{cases} \frac{t}{2} + 1 & \text{if } t \equiv 0 \pmod{2} \\ \frac{t+1}{2} & \text{if } t \equiv 1 \pmod{2} \end{cases}$


Thm [Ehrhart '62, Macdonald '71]

- if  $P$  has integral vertices then  $\text{ehr}_P(t)$  is polynomial in  $t$
- if  $P$  has rational vertices then  $\text{ehr}_P(t)$  is a quasi-polynomial
- if  $P$  is rational then
 
$$\text{ehr}_P(-t) = (-1)^{\dim P} \text{ehr}_{\text{Int}(P)}(t)$$

eg  $P = (0, \frac{1}{2}) \in \mathbb{R}$

$$\text{ehr}_P(t) = \begin{cases} \frac{t}{2} - 1 & \text{if } t \equiv 0 \pmod{2} \\ \frac{t-1}{2} & \text{if } t \equiv 1 \pmod{2} \end{cases}$$

FACT



"nice partition of  $Z_P$  into parallelepipeds"

$$[0, 2e_1) \times [0, 2e_2) + e_2$$

$$Z_P = Z_\emptyset \sqcup Z_1 \sqcup Z_2 \sqcup Z_3 \sqcup Z_{12} \sqcup Z_{13} \sqcup Z_{23}$$

$$\text{ehr}_P(t) = \text{vol}(P) t^{\dim P} \quad \text{if } P \text{ is an open parallelepiped}$$

ThM [Stanley '91, Ardila-Beck-McWhirter '20]

$\Gamma$  is a graph  $w \in \mathbb{R}^r$

$$\text{ehr}_{Z_\Gamma + w}(t) = \sum_{\substack{\mathcal{I} \text{ forest in } \Gamma \\ \text{Aff span}(Z_{\Gamma(\mathcal{I})} + tw) \cap Z^r \neq \emptyset}} \text{Vol}(\mathcal{I}) t^{|\mathcal{I}|}$$

$$\text{Aff span}(Z_{\Gamma(\mathcal{I})} + tw) \cap Z^r \neq \emptyset$$


DEF a flat  $T = \{T_1, T_2, \dots, T_k\} \vdash [r]$  is  $w$ -integral

if  $\forall \alpha \sum_{T \in T_\alpha} w_j \in \mathbb{Z}^{\text{span}(T_\alpha)} \quad w_\alpha \in \mathbb{R}$

Cor  $C(Z_{\Gamma} + w) = \sum_{\Gamma \text{ w-integral list}} (-1)^{r - l(\Gamma)} \sum_{F \text{ spanning forest of } \Gamma} \text{Vol}(F)$

Ex compute  $\#(\mathcal{X}^{-1}(a))$  for  $a \in A_n^{\text{red}}$  :

- $n=4 \quad d=0 \quad q=2$
- $n=4 \quad d=2 \quad q=2$



$\# \text{irr comp } \mathcal{X}^{-1}(a) = C(Z_{\Gamma_n} + w) \quad w_i = \frac{2n_i}{n}$

Intersection Cohomology

Q  $\text{IH}^*(M_{n,d}) = ?$  ↙ complex of sheaves

$\text{IH}^*(M_{n,d}; \mathbb{Q}) := H^*(M_{n,d}, \mathcal{IC})$   
 $= H^*(A_n, R\mathcal{X}_* \mathcal{IC})$

$R\mathcal{X}_* \mathcal{IC} |_{A_n^{\text{red}}}$

Theorem [Ngô, Macrì - Migliorini 2022]

$R\mathcal{X}_* \mathcal{IC} |_{\text{red}} \cong \bigoplus \mathcal{IC}_c(L_{n,d} \otimes \Lambda_n)$



where  $\cdot \Lambda_{\underline{n}}$  is the cohomology sheaf of the relative Picard group  $\text{Pic}^0(\bar{C}_{\underline{n}})$  of the normalization of  $C_{\underline{n}}$

$\cdot \mathcal{L}_{\underline{n},d}$  unknown coefficients  
 $\cdot$  local system on  $S_{\underline{n}}$

Q1  $\underline{n}, d \quad \mathcal{L}_{\underline{n},d} = 0$  ?

Q2  $\text{rk } \mathcal{L}_{\underline{n},d} = ?$

Q3 monodromy action

Remark  $\underline{a} \in S_{\underline{n}}$

$$\dim H^{\text{top}}(R\mathcal{X}_{\#} \mathcal{I}\mathcal{L})_{\underline{a}} = \# \text{ int. comp of } \mathcal{Z}^{-1}(\underline{a}) = C(Z_{\Gamma_{\underline{n}}} + w)$$

LEMMA  $a \in S_{\underline{n}} \quad \underline{n} = (n_1, \dots, n_r)$

$$H^{\text{top}}(R\mathcal{X}_{\#} \mathcal{I}\mathcal{L})_{\underline{a}} \cong \bigoplus_{\underline{I} + [\underline{r}]} (\mathcal{L}_{\underline{n}, \underline{I}, d})_{\underline{a}} \otimes \bigotimes_{i=1}^{\ell(\underline{I})} H^{\text{top}}(R\mathcal{X}_{n(\underline{I}), 0} \mathcal{I}\mathcal{L})_{\underline{a}}$$

$\uparrow$   $n, d$     $\uparrow$   $n(\underline{I}), 0$   
 $n(\underline{I}), d$

EX  $\underline{n} = \{1, 1, 2, 5\} \vdash 9 \quad r=4 \quad n=9$

$\underline{I} = \{13 | 24\} \vdash [4]$

$n(13) = 1+2 = 3$

$n(24) = 1+5 = 6$

$$(\mathcal{L}_{\{3,6\}, d})_{\underline{a}} \otimes H^{\text{top}}(R\mathcal{X}_{3,0} \mathcal{I}\mathcal{L})_{\{1,2\}} \otimes H^{\text{top}}(R\mathcal{X}_{6,0} \mathcal{I}\mathcal{L})_{\{1,2\}}$$

$$\in \mathcal{H}^{\text{top}}(\mathbb{R}\mathcal{X}_{g,d} \mathcal{L}\mathcal{C})_{\{1,1,2,3\}}$$

COR  $C(Z_{\Gamma_n} + w) = \sum_{T \in [\Gamma]} rK(L_{n_T, d}) C(Z_{\Gamma_T, \emptyset})$

$\forall \underline{n} \vdash n$

↑ restricted graph on the flat  $\underline{I}$

↑ solve this recurrence relation

$(n) \vdash n$   $C(Z_{\Gamma_n} + w) = rK(L_{(n), d}) C(Z_{\Gamma_n})$

$w \in \mathbb{Z}$

$\frac{d}{dn}$   $\underline{n} = (a, n-a)$   $(n) \rightsquigarrow rK(L_{(a, n-a), d})$

THM [Maun, Migliorini, P. 2023] For any graph  $w \in \mathbb{R}^+$


$$C(Z_{\Gamma} + w) = \sum_{S \in [\Gamma]} \left( \sum_{T \geq S} \mu_{\mathcal{L}}(S, T) \right) C(Z_{\Gamma_S})$$

COR  $rK(L_{\underline{n}, d}) = \sum_{\substack{T \in [\underline{n}] \\ T \text{ is } w\text{-integral}}} (-1)^{l(T)} \prod_{i=1}^{l(T)} (|T_i| - 1)!$

$\mu$  mobius function of the partition lattice



$L_{\Pi}$



$L_{\Pi, w}$

$\hat{0} = 1 | 2 | 3 | \dots | n$


$$\sum_{T \text{ w-integral}} \mu(\hat{\sigma}, T) = - \sum_{T \text{ non w-integral}} \mu(\hat{\sigma}, T) = \mu_{\mathcal{L}_{\Gamma, w}}(\hat{\sigma}, \hat{1})$$

$$\text{rk } \mathcal{L}_{n, d} = \mu_{\Gamma_n, w}(\hat{\sigma}, \hat{1}) = \tilde{\chi}(\Delta(\mathcal{L}_{\Gamma_n, w}))$$

THM Hall's Theorem

$$\mu_{\mathcal{L}}(\hat{\sigma}, \hat{1}) = \tilde{\chi}(\Delta(\mathcal{L} \setminus \{\hat{\sigma}, \hat{1}\}))$$

Shellability



LEX-shellability

good matching on chains in  $\mathcal{L}'' \rightsquigarrow$  prescribe a "good"

collapsing on  $\Delta(\mathcal{L} \setminus \{\hat{\sigma}, \hat{1}\})$

PROP if  $\mathcal{L}$  is LEX-shellable  $\Rightarrow \Delta(\mathcal{L} \setminus \{\hat{\sigma}, \hat{1}\}) \sim VS^{r-3}$

$$\mu_{\mathcal{L}}(\hat{\sigma}, \hat{1}) = (-1)^{r-3} \text{rk } H^{r-3}(\Delta(\mathcal{L} \setminus \{\hat{\sigma}, \hat{1}\}))$$


$$\text{if } n = (n) = \text{or } \frac{d \cdot n_i}{n} \notin \mathbb{Z} \Rightarrow \text{rk}(\mathcal{L}_{n, d}) > 0$$

THM [MacM, M, P]

$$(\mathcal{L}_{n, d})_p \cong H^{r-3}(\Delta(\mathcal{L}_{n, w} \setminus \{\hat{\sigma}, \hat{1}\})) \otimes \text{sgn}$$

as repr. of  $\pi_1(S_n) \rightarrow S_{k_1} \times S_{k_2} \times \dots \times S_{k_n}$   
 $\underline{n} = (1^{k_1}, 2^{k_2}, \dots, n^{k_n})$

1DBD compute characters



$M_{n,d}^{\text{red}}$