

# Poncelet's porism and periodic triangles in ellipse <sup>1</sup>

Vladimir Georgiev, Veneta Nedyalkova

## 1 Small historical introduction

One of the most important and beautiful theorems in projective geometry is that of Poncelet, concerning closed polygons which are inscribed in one conic and circumscribed about another (below we give the precise statement as well proof for the case of triangles). The theorem has deep interaction with other math fields. The aim of this section is to clarify one aspect of these relations: the connection between Poncelet's theorem and billiards in an ellipse. At first sight these topics seem unrelated, belonging to two distinct mathematical fields: geometry and dynamical systems. But there is a hidden thread tying these topics together: the existence of an underlying structure (we name it the Poncelet correspondence which turns out to be an elliptic curve. As is well known, elliptic curves can be endowed with a group structure, and the exploitation of this structure sheds much light on the aforementioned topics.

However, to read most of the books and available references some prerequisites (usually covered in undergraduate and first year graduate mathematics courses) are needed: complex analysis, linear algebra, and some point set topology.

In this sense the argument can not be adapted easily to some extracurricula activities in High Schools.

For this we are trying to find approach that needs only tools from the standard High School Programs.

This is not an easy problem. The classical A. Cayley (see [2], [3]) approach uses elliptic integrals, some other sources (see [5], [6], [8] and the references cited there) apply arguments for projective geometry and group theory.

The statement of the Poncelet's problem needs only to know the definition and the equation of the ellipse.

**Theorem 1.** (*Poncelet's Porism*) *Given one ellipse inside another, if there exists one circum-inscribed (simultaneously inscribed in the outer and circumscribed on the inner)  $n$ -gon, then any point on the boundary of the outer ellipse is the vertex of some circuminscribed  $n$ -gon.*

There are several proofs of this remarkable theorem, most of which are not elementary. Poncelet's theorem dates to the nineteenth century and has attracted the attention of many mathematicians of that period (a detailed historical account is given in [1]). The main reason for this interest seems to stem from the fact that several proofs of this theorem require the use of complex and homogeneous coordinates, notions which were beginning to emerge at the time (1813) when Poncelet discovered his theorem. Poncelet discovered the theorem while in captivity as war

---

<sup>1</sup>This project has been funded with support from the European Commission in its Lifelong Learning Programme (510028-LLP-1-2010-1-IT-COMENIUS-CMP). This publication reflects the views only of the authors, and the Commission cannot be held responsible for any use which may be made of the information contained therein




Figure 1: Poncelet's theorem for the case of circle and ellipse.

prisoner in the Russian city of Saratov. After his return to France, a proof appears in his book [7], published in 1822. The proof, which is synthetic and somewhat elaborate, reduces the theorem to two (not' necessarily concentric) circles. A discussion of the ideas in Poncelet's proof is given in [1], pp. 298-311.

Our purpose is to find elementary proof in one nontrivial situation: the case  $n = 3$  and the situation, when we have two ellipses

$$e : \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad (1)$$

and

$$e_1 : \frac{x^2}{a_1^2} + \frac{y^2}{b_1^2} = 1, \quad (2)$$

such that  $e_1$  is inside  $e$ .

We shall prove in this case the Poncelet' theorem as well as the following more precise result.

**Theorem 2.** (see Figure 1 ) Suppose the ellipse (2) is inside the ellipse (1), i.e.

$$a > b > 0, a_1 > b_1 > 0,$$

$$a > a_1, b > b_1.$$

Then the following conditions are equivalent:

i) there exists a triangle  $\triangle A_0B_0C_0$  inscribed in  $e$  and circumscribed on  $e_1$ ,

ii) we have the relation

$$\frac{a_1}{a} + \frac{b_1}{b} = 1.$$

iii) for any point  $A$  on the ellipse  $e$  one can find a unique triangle  $\triangle ABC$  inscribed in  $e$  and circumscribed on  $e_1$ .

## 2 Reduction to the case of circle and ellipse and preliminary facts

Consider two ellipses

$$e : \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad (3)$$

and

$$e_1 : \frac{x^2}{a_1^2} + \frac{y^2}{b_1^2} = 1, \quad (4)$$

such that  $e_1$  is inside  $e$ . This condition can be expressed as

$$a > b > 0, a_1 > b_1 > 0,$$

$$a > a_1, b > b_1.$$

One can use a simple change of coordinates in the plane

$$X = \frac{x}{a}, \quad Y = \frac{y}{b}, \quad (5)$$

so that the ellipse  $e$  in the new coordinates  $X, Y$  has equation

$$X^2 + Y^2 = 1. \quad (6)$$

so it is the circle  $k(O, 1)$  with center at the origin  $O$  of the new coordinate system and has radius 1.

The second ellipse  $e_1$  becomes

$$\frac{X^2}{A_1^2} + \frac{Y^2}{B_1^2} = 1, \quad A_1 = \frac{a_1}{a}, B_1 = \frac{b_1}{b} \quad (7)$$

and it is clear that this change of coordinates preserves the notions of intersection, line is transformed in line, circle in circle, ellipse in ellipse (or circle as a partial case) and if the line and ellipse are tangent they remain tangent after the change of the coordinates (see Figure 2).

**Exercise 1.** *Prove the fact that if line and ellipse are tangent they remain tangent after the change of the coordinates (5).*




Figure 2: Ellipse is transformed in circle.

For this from now on we shall work with circle  $k(O, 1)$  with center at the origin  $O$  and radius 1

$$x^2 + y^2 = 1. \quad (8)$$

and ellipse  $e_1$

$$\frac{x^2}{a_1^2} + \frac{y^2}{b_1^2} = 1, \quad 1 > a_1 \geq b_1 \quad (9)$$

inside  $k(O, 1)$  as it is shown on Figure 1.

We prepare again a list of questions preparing the solution of the problem (or proof of the Poncelet's theorem):

- Given an ellipse  $e_1 : x^2/a_1^2 + y^2/b_1^2 = 1$  and the point  $A_0(x_0, y_0)$  on  $k(O, 1)$  find the tangent lines from  $A_0$  to  $e_1$  and find also the points  $A_1, A_2$  of the intersection of these tangent lines with the circle  $x^2 + y^2 = 1$  (we need formula expressing the coordinates of  $A_1, A_2$  in terms of  $x_0, y_0$  and the angular coefficients  $k_1, k_2$  of the lines  $A_0A_1$  and  $A_0A_2$  respectively;
- Using the parametrization

$$x_j = \cos \varphi_j, y_j = \sin \varphi_j, \quad j = 0, 1, 2 \quad (10)$$

find a relation between  $\varphi_j$  and  $\theta_{1,2} = \arctan k_{1,2}$ .

- Given an ellipse  $e_1 : x^2/a_1^2 + y^2/b_1^2 = 1$ , the point  $A_0(x_0, y_0)$  on  $k(O, 1)$ , the tangent lines from  $A_0$  to  $e_1$  intersecting  $k(O, 1)$  into the points  $A_1, A_2$  and using the parametrization (10) express the necessary and sufficient condition that the line  $A_0A_1$  is tangent to the ellipse  $e_1$  in terms of  $\varphi_0, \varphi_1$  and  $\theta_1 = \arctan k_1$ .
- Given an ellipse  $e_1 : x^2/a_1^2 + y^2/b_1^2 = 1$ , the point  $A_0(x_0, y_0)$  on  $k(O, 1)$ , the tangent lines from  $A_0$  to  $e_1$  intersecting  $k(O, 1)$  into the points  $A_1, A_2$  and using the parametrization (10) express the necessary and sufficient condition that the line  $A_0A_2$  is tangent to the ellipse  $e_1$  in terms of  $\varphi_0, \varphi_2$  and  $\theta_2 = \arctan k_2$ .
- Using simple trigonometric transformations show that the following two conditions a) the line  $A_0A_1$  is tangent to the ellipse  $e_1$  (condition is expressed in terms of  $\varphi_0, \varphi_1$  and  $\theta_1 = \arctan k_1$ ) b) the line  $A_0A_2$  is tangent to the ellipse  $e_1$  (condition is expressed in terms of  $\varphi_0, \varphi_2$  and  $\theta_2 = \arctan k_2$ ) imply a) the line  $A_1A_2$  is tangent to the ellipse  $e_1$  (condition is expressed in terms of  $\varphi_1, \varphi_2$  and  $\theta_{1,2} = \arctan k_{1,2}$ )

Step by step we give answers presenting some Lemmas that can be verified without difficulty.

**Lemma 1.** *Given an ellipse  $e_1 : x^2/a_1^2 + y^2/b_1^2 = 1$  one can express the necessary and sufficient condition such that the line  $y - y_0 = k(x - x_0)$  through the point  $A_0(x_0, y_0)$  is tangent to  $e_1$  as follows*

$$(y_0 - kx_0)^2 = b_1^2 + k^2 a_1^2.$$

**Lemma 2.** *Given an ellipse  $e_1 : x^2/a_1^2 + y^2/b_1^2 = 1$  and point  $A_0(x_0, y_0)$  on the unit circle and denote by*

$$t : y - y_0 = k(x - x_0)$$

*any line through  $A_0$  and by  $A_1(x_1, y_1)$  the point of the second intersection of this line with the unit circle  $k(O, 1) : x^2 + y^2 = 1$ , such we have*

$$x_1 = \frac{k^2 - 1}{k^2 + 1}x_0 - \frac{2k}{k^2 + 1}y_0,$$

$$y_1 = -\frac{2k}{k^2 + 1}x_0 - \frac{k^2 - 1}{k^2 + 1}y_0.$$




Figure 3: When  $A_0A_1$  is tangent to  $e_1$ ?

*Proof.* The intersection points are given by the equations

$$x^2 + (y_0 + k(x - x_0))^2 = 1.$$

This equation has two roots  $x_0$  and  $x_1$  so

$$x_0 + x_1 = -\frac{2k(y_0 - kx_0)}{1 + k^2}.$$

From this relation we get the expression for  $x_1$ . Similarly we proceed for  $y_1$ .  $\square$

**Lemma 3.** Given an ellipse  $e_1 : x^2/a_1^2 + y^2/b_1^2 = 1$  and a point  $A_0(\cos \varphi_0, \sin \varphi_0)$  on the unit circle denote by

$$t : y - y_0 = k(x - x_0)$$

any line from  $A_0$  and let  $A_1$  the second point of intersection of this lines with the circle  $k(O, 1) : x^2 + y^2 = 1$ , such that  $A_1(\cos \varphi, \sin \varphi)$ . Then the relations of Lemma 2 take the form we have

$$\theta = \frac{\varphi + \varphi_0 - \pi}{2} + m\pi, m \in \mathbb{Z},$$

where

$$\theta = \arctan k.$$

*Proof.* We have the relations

$$\frac{k^2 - 1}{k^2 + 1} = -\cos(2\theta), \quad \frac{2k}{k^2 + 1} = \sin(2\theta).$$

Making the substitution

$$x_1 = \cos \varphi, y_1 = \sin \varphi$$

we find

$$\cos \varphi = -\cos(2\theta) \cos \varphi_0 - \sin(2\theta) \sin \varphi_0 =$$

$$\begin{aligned}
&= \cos(2\theta + \pi) \cos \varphi_0 + \sin(2\theta + \pi) \sin \varphi_0 = \cos(2\theta + \pi - \varphi_0), \\
&\sin \varphi = -\sin(2\theta) \cos \varphi_0 + \cos(2\theta) \sin \varphi_0 = \\
&= \sin(2\theta + \pi) \cos \varphi_0 - \cos(2\theta + \pi) \sin \varphi_0 = \sin(2\theta + \pi - \varphi_0),
\end{aligned}$$

and these relations lead simply to the needed relation

$$2\theta + \pi - \varphi_0 = \varphi + 2m\pi, m \in \mathbb{Z}.$$

This completes the proof. □

**Lemma 4.** *Given an ellipse  $e_1 : x^2/a_1^2 + y^2/b_1^2 = 1$  and a point  $A_0(\cos \varphi_0, \sin \varphi_0)$  denote by*

$$t : y - y_0 = k(x - x_0)$$

*a line through  $A_0$  and by  $A_1$  the point of the intersection of this line with the circle  $e : x^2 + y^2 = 1$ , such that  $A_1(\cos \varphi, \sin \varphi)$ . Then  $t$  is tangent to  $e_1$  if and only if*

*we have*

$$\cos^2 \left( \frac{\varphi - \varphi_0}{2} \right) = b_1^2 \sin^2 \left( \frac{\varphi + \varphi_0}{2} \right) + a_1^2 \cos^2 \left( \frac{\varphi + \varphi_0}{2} \right) = (a_1^2 - b_1^2) \cos^2 \left( \frac{\varphi + \varphi_0}{2} \right) + b_1^2.$$

*Proof.* From Lemma 1 we see that we need to transform  $(y_0 - kx_0)^2$  into a function of  $\varphi$  and  $\varphi_0$ . Indeed, we have

$$y_0 - kx_0 = \frac{\cos \theta \sin \varphi_0 - \sin \theta \cos \varphi_0}{\cos \theta} = \frac{\sin(\varphi_0 - \theta)}{\cos \theta}. \quad (11)$$

Using now the relation

$$\theta = \frac{\varphi + \varphi_0 - \pi}{2} + m\pi, m \in \mathbb{Z},$$

from Lemma 1, we see the the numerator in (11) is

$$\sin(\varphi_0 - \theta) = \sin \left( \frac{\varphi_0 - \varphi + \pi}{2} - m\pi \right) = (-1)^m \cos \left( \frac{\varphi_0 - \varphi}{2} \right)$$

while the denominator becomes

$$\cos \theta = \cos \left( \frac{\varphi + \varphi_0 - \pi}{2} + m\pi \right) = (-1)^m \sin \left( \frac{\varphi + \varphi_0}{2} \right)$$

so we find

$$\sin^2 \left( \frac{\varphi - \varphi_0}{2} \right) (y_0 - kx_0)^2 = \cos^2 \left( \frac{\varphi - \varphi_0}{2} \right).$$

Applying Lemma 1 combined with the above relations, we complete the proof of the Lemma. □

**Remark 1.** *We can rewrite the relations of Lemma 4 in different ways using the formula*

$$\cos^2 \alpha = \frac{1 + \cos(2\alpha)}{2},$$

*also as*

$$\cos(\varphi - \varphi_0) = c^2 \cos(\varphi + \varphi_0) + D, \quad (12)$$

*or*

$$(1 - c^2) \cos \varphi \cos \varphi_0 + (1 + c^2) \sin \varphi \sin \varphi_0 = D, \quad (13)$$

*where*

$$c^2 = a_1^2 - b_1^2, D = a_1^2 + b_1^2 - 1. \quad (14)$$

### 3 Proof of Poncelet theorem using trigonometric functions




Figure 4: The meaning of the assumption  $\triangle A_0B_0C_0$  is circumscribed on  $e_1$ ?

We take a point  $A_0(\cos \varphi_0, \sin \varphi_0)$  on the unit circle and find of two tangent lines  $t_1, t_2$  through  $A_0$  to the ellipse

$$e_1 = \frac{x^2}{a_1^2} + \frac{y^2}{b_1^2} = 1.$$

Then we find the intersection points of  $t_1, t_2$  with the unit circle (see Figure 4) and denote the two intersection points (different from  $A_0$ ) by

$$B_0(\cos \varphi_1, \sin \varphi_1), C_0(\cos \varphi_2, \sin \varphi_2).$$

First, let us express the assumption of Poncelet's theorem that there exists at least one triangle  $\triangle A_0B_0C_0$  inscribed in the unit circle, i.e.

$$A_0(\cos \varphi_0, \sin \varphi_0), B_0(\cos \varphi_1, \sin \varphi_1), C_0(\cos \varphi_2, \sin \varphi_2), \quad 0 \leq \varphi_0 < \varphi_1 < \varphi_2 \leq 2\pi$$

and circumscribed on the inner ellipse  $e_1$ . Since  $A_0B_0$  is tangent to  $e_1$  we know that:

$$\cos^2 \left( \frac{\varphi_1 - \varphi_0}{2} \right) = (a_1^2 - b_1^2) \cos^2 \left( \frac{\varphi_1 + \varphi_0}{2} \right) + b_1^2 \quad (15)$$

(this is due to Lemma 4). Similarly, the fact that  $A_0C_0$  and  $B_0C_0$  are tangent to  $e_1$ , and Lemma 4 imply

$$\cos^2 \left( \frac{\varphi_2 - \varphi_0}{2} \right) = (a_1^2 - b_1^2) \cos^2 \left( \frac{\varphi_2 + \varphi_0}{2} \right) + b_1^2. \quad (16)$$

$$\cos^2 \left( \frac{\varphi_2 - \varphi_1}{2} \right) = (a_1^2 - b_1^2) \cos^2 \left( \frac{\varphi_2 + \varphi_1}{2} \right) + b_1^2. \quad (17)$$

We can unify all these relations into one

$$\cos^2 \left( \frac{\varphi_j - \varphi_\ell}{2} \right) = (a_1^2 - b_1^2) \cos^2 \left( \frac{\varphi_j + \varphi_\ell}{2} \right) + b_1^2, \quad 0 \leq j \neq \ell \leq 2. \quad (18)$$

Take any point  $A(\cos\psi_0, \sin\psi_0)$  on the unit circle and find of two tangent lines  $t_1, t_2$  through  $A_0$  to the ellipse

$$e_1 : \frac{x^2}{a_1^2} + \frac{y^2}{b_1^2} = 1.$$

Then we find the intersection points of  $t_1, t_2$  with the unit circle (see Figure 5) and denote the two intersection points (different from  $A$ ) by

$$B(\cos\psi_1, \sin\psi_1), C(\cos\psi_2, \sin\psi_2).$$




Figure 5: Two sides tangent  $\Rightarrow$  the third side is also tangent.

Since  $AB$  is tangent to  $e_1$  we know that:

$$\cos^2\left(\frac{\psi_1 - \psi_0}{2}\right) = (a_1^2 - b_1^2) \cos^2\left(\frac{\psi_1 + \psi_0}{2}\right) + b_1^2 \quad (19)$$

(this is due to Lemma 4). Similarly, the fact that  $A_0C_0$  and  $B_0C_0$  are tangent to  $e_1$ , and Lemma 4 imply

$$\cos^2\left(\frac{\psi_2 - \psi_0}{2}\right) + (a_1^2 - b_1^2) \cos^2\left(\frac{\psi_2 + \psi_0}{2}\right) = b_1^2. \quad (20)$$

So we summarize all assumptions of Poncelet's theorem and can say that (18), (19) and (20) are satisfied.

What we have to prove?

Having in mind again Lemma 4 we see that our purpose is to show that

$$\cos^2\left(\frac{\psi_2 - \psi_1}{2}\right) = (a_1^2 - b_1^2) \cos^2\left(\frac{\psi_2 + \psi_1}{2}\right) + b_1^2. \quad (21)$$



This relation can be rewritten as

$$(1 - c^2) \cos \psi_2 \cos \psi_1 = (1 + c^2) \sin \psi_2 \sin \psi_1 + D, \quad (22)$$

where

$$c^2 = a_1^2 - b_1^2, D = a_1^2 + b_1^2 - 1. \quad (23)$$

according to Remark 1.

Now we are in position to apply the trigonometric lemma from the appendix and conclude that

$$\cos^2 \left( \frac{\psi_2 - \psi_1}{2} \right) = \frac{4c^2 D^2}{(1 - c^2)^2 (1 + c^2)^2} \cos^2 \left( \frac{\psi_2 + \psi_1}{2} \right) + \frac{D^2}{(1 + c^2)^2}. \quad (24)$$

Comparing this relation with (21) we see that the following conditions

$$4D^2 = (1 - c^2)^2 (1 + c^2)^2, \quad D^2 = b_1^2 (1 + c^2)^2 \quad (25)$$

are required. This relations and (23) lead to the following sufficient condition

$$a_1 + b_1 = 1 \quad (26)$$

that implies  $\triangle ABC$  is circumscribed on  $e_1$ . The condition (23) is also necessary for the fulfillment of the property

- there exists a triangle  $\triangle A_0 B_0 C_0$  circumscribed on  $e_1$ .

If there exists at least one  $\triangle A_0 B_0 C_0$  circumscribed on  $e_1$ , then (26) and hence  $\triangle ABC$  is circumscribed on  $e_1$ .

This completes the proof of the Theorem.

## 4 Appendix: Trigonometric Lemma

**Lemma 5.** *Suppose*

$$\sin \left( \frac{\psi_1 - \psi_2}{2} \right) \neq 0, \quad \cos \left( \frac{\psi_1 + \psi_2}{2} \right) \neq 0, \quad \cos \psi_0$$

and

$$\begin{cases} (1 - c^2) \cos \psi_1 \cos \psi_0 + (1 + c^2) \sin \psi_1 \sin \psi_0 = D & ; \\ (1 - c^2) \cos \psi_2 \cos \psi_0 + (1 + c^2) \sin \psi_2 \sin \psi_0 = D & . \end{cases} \quad (27)$$

Then

$$(1 - c^2) \tan \left( \frac{\psi_1 + \psi_2}{2} \right) = (1 + c^2) \tan \psi_0 \quad (28)$$

and moreover

$$\cos^2 \left( \frac{\psi_2 - \psi_1}{2} \right) = \frac{4c^2 D^2}{(1 - c^2)^2 (1 + c^2)^2} \cos^2 \left( \frac{\psi_2 + \psi_1}{2} \right) + \frac{D^2}{(1 + c^2)^2}. \quad (29)$$

*Proof.* Take the difference between the relations in (27). We get

$$-(1-c^2) \sin\left(\frac{\psi_1 - \psi_2}{2}\right) \sin\left(\frac{\psi_1 + \psi_2}{2}\right) \cos \psi_0 + (1+c^2) \sin\left(\frac{\psi_1 - \psi_2}{2}\right) \cos\left(\frac{\psi_1 + \psi_2}{2}\right) \sin \psi_0 = 0.$$

The assumption

$$\sin\left(\frac{\psi_1 - \psi_2}{2}\right) \neq 0$$

implies that

$$(1 - c^2) \sin\left(\frac{\psi_1 + \psi_2}{2}\right) \cos \psi_0 = (1 + c^2) \cos\left(\frac{\psi_1 + \psi_2}{2}\right) \sin \psi_0.$$

This proves (28). The other relation can be obtained following the plan

- first equation in (27)  $\times \sin \psi_2$  – second equation in (27)  $\times \sin \psi_1$ ;
- first equation in (27)  $\times \cos \psi_2$  – second equation in (27)  $\times \cos \psi_1$ .

In this way we get

$$\begin{aligned} 2D \sin\left(\frac{\psi_2 - \psi_1}{2}\right) \cos\left(\frac{\psi_2 + \psi_1}{2}\right) &= 2(1 - c^2) \sin\left(\frac{\psi_2 - \psi_1}{2}\right) \cos\left(\frac{\psi_2 + \psi_1}{2}\right) \cos \psi_0, \\ -2D \sin\left(\frac{\psi_2 - \psi_1}{2}\right) \sin\left(\frac{\psi_2 + \psi_1}{2}\right) &= -2(1 + c^2) \sin\left(\frac{\psi_2 - \psi_1}{2}\right) \cos\left(\frac{\psi_2 + \psi_1}{2}\right) \sin \psi_0, \end{aligned}$$

so using the assumption

$$\sin\left(\frac{\psi_1 - \psi_2}{2}\right) \neq 0$$

we find

$$\begin{aligned} \frac{D}{1 - c^2} \cos\left(\frac{\psi_2 + \psi_1}{2}\right) &= \cos\left(\frac{\psi_2 - \psi_1}{2}\right) \cos \psi_0, \\ \frac{D}{1 + c^2} \sin\left(\frac{\psi_2 + \psi_1}{2}\right) &= \cos\left(\frac{\psi_2 - \psi_1}{2}\right) \sin \psi_0. \end{aligned}$$

Taking the sum of squares of these identities we obtain

$$\frac{D^2}{(1 - c^2)^2} \cos^2\left(\frac{\psi_2 + \psi_1}{2}\right) + \frac{D^2}{(1 + c^2)^2} \sin^2\left(\frac{\psi_2 + \psi_1}{2}\right) = \cos^2\left(\frac{\psi_2 - \psi_1}{2}\right)$$

and this equation yields (29).

This completes the proof of the Lemma.

□

## References

- [1] H. J. M. Bos, C. Kers, F. Oort, and D. W. Raven, *Poncelet's closure theorem*, Expo. Math. **5** (1987), 289 – 364.
- [2] A. Cayley, *Note on the porism of the in-and-circumscribed polygon*, Philosophical magazine **6** (1853), 99102.
- [3] A. Cayley, *Developments on the porism of the in-and-circumscribed polygon*, Philosophical magazine **7** (1854), 339345.
- [4] H. Dörrie, *100 great problems in Mathematics. Their history and solutions*, Dover Publ., New York, (1965).
- [5] V. Dragovic, M. Radnovic *Poncelet Porisms and Beyond*, Birkhäuser, Springer-Basel, (2011).
- [6] L.Flatto, *Poncelet's Theorem*, AMS, (2008).
- [7] J. V. Poncelet, *Traite sur les Proprietes des Figures*, Paris, (1822).
- [8] S.Tabachnikov, *Geometry and Billiards*, Students Mathematica Library, (2005)